

**EFFECT OF A DIET RICH IN OLIVE OIL PHENOLICS ON AGE-RELATED
CHANGES IN MOUSE MODEL OF AGEING AND HPLC ANALYSIS OF OLIVE OIL
PHENOLICS WITH ELECTROCHEMICAL DETECTOR**

**Ph.D. Thesis by
Banu BAYRAM**

Department : Food Engineering

Programme : Food Engineering

SEPTEMBER 2011

**EFFECT OF A DIET RICH IN OLIVE OIL PHENOLICS ON AGE-RELATED
CHANGES IN MOUSE MODEL OF AGEING AND HPLC ANALYSIS OF OLIVE OIL
PHENOLICS WITH ELECTROCHEMICAL DETECTOR**

**Ph.D. Thesis by
Banu BAYRAM
(506052501)**

Date of submission : 6 September

Date of defence examination : 29 September

Supervisor (Chairman) : Assoc. Prof. Dr. Beraat OZCELIK (ITU)
Co-Advisor : Prof. Dr. Gerald RIMBACH (CAU)
Members of the Examining Committee : Prof. Dr. Funda ELMACIOGLU (MU)
Prof. Dr. A. Süha YALCIN (MU)
Prof. Dr. Emine UBAY COKGOR (ITU)
Assoc. Prof. Dr. Gürbüz GUNES (ITU)
Assoc. Prof. Dr. Hakan BERMEK (ITU)

SEPTEMBER 2011

**ZEYTİNYAĞI FENOLİKLERİNCE ZENGİN DİYETİN YAŞLANMASI
HIZLANDIRILMIŞ FARE MODELİNDE YAŞLANMAYA BAĞLI DEĞİŞİKLİKLER
ÜZERİNE ETKİSİ VE ZEYTİNYAĞINDAKİ FENOLİK MADDELERİN
ELEKTROKİMYASAL DEDEKTÖR İLE HPLC ANALİZİ**

**DOKTORA TEZİ
Banu BAYRAM
(506052501)**

Tezin Enstitüye Verildiği Tarih : 06 Eylül 2011

Tezin Savunulduğu Tarih : 29 Eylül 2011

**Tez Danışmanı : Doç. Dr. Beraat ÖZÇELİK (İTÜ)
Eş Danışmanı : Prof. Dr. Gerald RIMBACH (CAÜ)
Diğer Jüri Üyeleri : Prof. Dr. Funda ELMACIOĞLU (MÜ)
Prof. Dr. A.Süha YALÇIN (MÜ)
Prof. Dr. Emine UBAY ÇOKGÖR (İTÜ)
Doç. Dr. Gürbüz GÜNEŞ (İTÜ)
Doç. Dr. Hakan BERMEK (İTÜ)**

EYLÜL 2011

FOREWORD

Foremost, I would like to thank to Assoc. Prof. Dr. Beraat ÖZCELIK for her continual support, trust and love throughout this study. Her advice, encouraging and personal guidance have been of great value for me. I am proud of being her “daughter”.

I would like to express my special thanks to Prof. Dr Gerald RIMBACH for his continual support, advice, knowledge and expertise, understanding, detailed and constructive comments, and for his kind support throughout this work.

I partly performed my PhD studies in Germany and collaborated with many colleagues. I wish to express my sincere thanks to all those who have helped me with my work at Christian Albrechts University of Kiel Institute of Human Nutrition and Food Science. Firstly I would like to give my special thanks to Dr. Jan FRANK, who helped me during my thesis and dealt all the problems that I faced. Many thanks for his positive energy, support and always having time for me. I would like to thank to Dr. Sonja FRANK for her friendship and support. I would like to thank to my very special friend Dr. Christoph HUNDHAUSEN for his great support. I would like to thank to Dr. Chi Vinh Duong from Phenomenex for his kind support in HPLC studies.

Also I would like to thank to my friends in Kiel, Dr. Tuba ESATBEYOGLU, Dr. Insa ERNST, Anne GREASER, Charlotte SCHRADER, Gaby STEINKAMP, Anke SCHLÖSSER, Christine STURM, Jennifer MÖHRING and Janina DOSE for their collaboration and friendships.

During my PhD studies I was supported by TUBITAK (The Scientific and Technological Research Council of Turkey) and I would like to thank to this council for the financial support.

I wish to thank to my family, for all love and continual support on the this long way till getting my PhD title. I dedicate this work to my mother and father. I hope she is proud of me.

September 2011

Banu Bayram

Biologist

TABLE OF CONTENTS

	<u>Page</u>
FOREWORD	v
TABLE OF CONTENTS	vii
ABBREVIATIONS	xi
LIST OF TABLES	xiii
LIST OF FIGURES	xv
SUMMARY	xvii
ÖZET	xxi
1. INTRODUCTION	1
2. LITERATURE	5
2.1 Production of Olive Oil	5
2.2 Chemical Composition of Olive Oil	6
2.2.1 Major compounds	6
2.2.2 Minor compounds	6
2.2.2.1 Lipophilic phenols (Vitamin E)	7
2.2.2.2 Sterols	7
2.2.2.3 Hydrocarbons	7
2.2.2.4 Pigments	7
2.2.2.5 Volatile compounds	7
2.2.2.6 Triterpenes	8
2.3 Health Benefits of Olive Oil	8
2.4 Hydrophilic Phenolic Compounds	10
2.5 Factors that Affect the Phenolic Content of Olive Oil	12
2.5.1 Agronomic conditions	12
2.5.2 Technological conditions	13
2.6 Health Effects of Olive Oil Phneolics	14
2.7 Analytical Determination of Phenolic Compounds in Olive Oil	18
2.8 Principle of the Electrochemical Detector	19
2.9 Advantages of Electrochemicaal Detector	19
2.10 Mechanism of Ageing and Ageing Related Theories	20
2.11 Biomarkers of Oxidative Stress	22
2.11.1 Lipid peroxidation	23
2.11.2 Nucleic acid oxidation	23
2.11.3 Protein oxidation	23
2.11.4 Decreased proteasomal activity	24
2.11.5 Increased inflammation	24
2.12 Protection from Detrimental Effects of Oxidative Stress	25
2.12.1 Antioxidant defence system	25
2.12.1.1 Vitamin C (Ascorbic Acid)	25
2.12.1.2 Vitamin E (α -Tocopherol)	25
2.12.1.3 Glutathione (GSH)	26
2.12.1.4 Uric acid	27
2.12.1.5 Paraoxonase (PON)	28
2.12.2 Phase II metabolizing enzymes	29
2.12.2.1 NADPH quinone oxidoreductase (NOQ-1)	29

2.12.2.2 Glutathione-S-transferase (GST)	29
2.12.2.3 Gamma glutamyl cysteinyl synthetase (γ -GCS)	20
2.12.2.4 Hemeoxygenase-1 (HO-1)	30
2.13 Models of Human Ageing and the Senescence Accelerated Mouse Prone (SAMP)	31
2.14 In Vivo Studies Related to Effects of Olive Oil on Ageing	32
2.15 Aim of the Study	33
3. MATERIALS AND METHODS	35
3.1 Chemicals and Reagents	35
3.2 Sampling of Olive Oil	35
3.3 Preparation of Olive Oil Extracts	36
3.4 Spectrophotometric Determination of Total Phenols	36
3.5 HPLC Analysis of Olive Oil Phenolics	36
3.6 Method Validation	37
3.6.1 Selectivity	37
3.6.2 Linearity lower limit of detection (LLOD) and lower limit of quantification (LLOQ)	37
3.6.3 Accuracy	38
3.6.4 Precision	38
3.6.5 Recovery	38
3.6.6 Stock solution stability	38
3.6.7 Short term stability	38
3.6.8 Freeze thaw stability	39
3.6.9 Post preparative stability	39
3.7 FRAP and TEAC Assays	39
3.8 Animals and Study Design	39
3.9 PON1 Arylesterase Activity in Serum	40
3.10 Liver and Heart Tissue Preparation	40
3.11 Lipid Peroxidation	40
3.12 Quantification of Protein Carbonyls	40
3.13 Proteasomal Activity	41
3.14 α -Tocopherol Analysis	41
3.15 Quantification of Ascorbic Acid, GSH, GSSG, Uric Acid, Cysteine, Homocysteine and Methionine by HPLC with Coulometric Electrochemical Detection	42
3.16 RNA Isolation and Real Time Quantitative RT-PCR	42
3.17 Western Blot Analysis	44
3.18 Statistical Analysis	45
4. RESULTS AND DISSCUSSION	47
4.1 Method Validation	47
4.1.1 Selectivity	47
4.1.2 Linearity lower limit of detection (LLOD) and lower limit of quantification (LLOQ)	47
4.1.3 Accuracy, precision and recovery	48
4.1.4 Stability studies	50
4.2 Application of the Method for The Determinaton of Selected Phenolic Compounds in Olive Oils	53
4.3 Animal Study	56
4.3.1 Body weight and feed intake	56
4.3.2 Effect of age and HP diet on serum parameters	57
4.3.3 Effect of age and HP diet on on oxidative stress biomarkers	58
4.3.4 Effect of age and HP diet on redox status	60
4.3.5 Effect of age and HP diet on antioxidant vitamins in liver	62
4.3.6 Effect of age and HP diet on antioxidant phase II enzymes	67
4.3.7 Effect of age and HP diet on SIRT1 expression	71

4.3.8 Effect of age and HP diet on inflammation markers	72
5. CONCLUSION	75
REFERENCES	79
APPENDICES	103
CURRICULUM VITAE	113

ABBREVIATIONS

ABCA1	: ATP binding cassette A1
CRP	: C reactive protein
CYP3A	: Cytochrome P450 family 3A
EVOO	: Extra virgin olive oil
GAPDH	: Glyceraldehyde phosphate dehydrogenase
GSH	: Glutathion
GSSG	: Glutathion disulphite
GST	: Glutathion-S-transferase
γ-GCS	: γ-glutamyl-cysteinyl-synthetase
HMGCoAR	: 3-hydroxy-3-methyl-glutaryl-CoA reductase
HO-1	: Hemeoxygenase-1
HP	: High phenolic
LCAT	: Lecithin Cholesterin Acyltransferase
LDLrec	: Low density lipoprotein receptor
LLE	: Liquid liquid extraction
LLOQ	: Lower limit of quantification
LLOD	: Lower limit of detection
LP	: Low phenolic
LPL	: Lipoprotein lipase
LRP-1	: Low density lipoprotein receptor related protein 1
MDR-1	: Multidrug resistance transporter 1
NQO-1	: NADPH quionone oxidoreductase
Nrf2	: Nuclear factor erythroid 2-related factor 2
oxo8dG	: 8-oxo-2-deoxyguanosine
PON-1	: Paraoxonase-1
PON-2	: Paraoxonase-2
SAMP8	: Senescence Accelerated Mouse Prone 8
SAMR1	: Senescence Accelerated Mouse Resistant 1
SAP	: Serum amyloid P
SIRT1	: Sirtuin1
SPE	: Solid phase extraction
SRB1	: Scavenger receptor B1
SVCT1	: Na- dependent vitamin C transporter-1
SVCT2	: Na- dependent vitamin C transporter-2
TBARS	: Thiobarbituric acid reactive substances
α-TTP	: Alpha tocopherol transfer protein
VOO	: Virgin olive oil

LIST OF TABLES

	<u>Page</u>
Table 2.1 : Chemical composition of olive oil	6
Table 2.2 : Mechanism of actions of olive oil components in their health effects ...	9
Table 2.3 : Phenolic compounds in olive oil	11
Table 2.4 : Health effect of the main olive oil phenolic compounds	16
Table 3.1 : Primers used in real time PCR experiment	43
Table 4.1 : Linear range, LLOQ, LLOD of selected phenolic compounds for the method validation	48
Table 4.2 : Intraday and inter-day recoveries, accuracies and precisions of selected phenolic compounds of olive oil	49
Table 4.3 : Stock solution stability of standards after 24 h storage at room temperature, 4°C and -20°C	51
Table 4.4 : Freeze thaw stability and short term stability of selected phenolic compounds in olive oil at room temperature, 4°C, -20°C, and 80°C .	52
Table 4.5 : Post preparative stability of selected phenolic compounds in olive oil	53
Table 4.6 : FRAP, TEAC, total phenolic, alpha tocopherol, gamma tocopherol and phenolic contents of collected olive oil samples	54
Table 4.7 : Concentrations of phenols and TEAC and FRAP value of the High-Polyphenol (HP) and Low-Polyphenol (LP) olive oils.....	56
Table 4.8 : Body weight and food intake of SAMR1 and SAMP8 mice fed either with high or low phenolic diet	57
Table 4.9 : Serum parameters of SAMR1 and SAMP8 mice	58
Table 4.10 : Concentrations of TBARS and protein carbonyls as well as proteasomal activities in liver homogenates from SAMR1 and SAMP8 mice	59
Table 4.11 : Concentrations of TBARS and protein carbonyls as well as proteasomal activities in heart homogenates from SAMR1 and SAMP8 mice	59
Table 4.12 : The concentration of GSH, GSSG, GSH/GSSG, cysteine, methionine and uric acid in liver of SAMR1 and SAMP8 mice	61
Table 4.13 : The concentration of GSH, GSSG, GSH/GSSG, cysteine, methionine and uric acid in heart of SAMR1 and SAMP8 mice	61
Table 4.14 : Relative mRNA expression (normalized for GAPDH) of SVCT1, SVCT2 and Gulonolactone oxidase in liver of SAMR1 and SAMP8 mice	64
Table C.1 : The list of the analysed olive oil samples	109
Table D.1 : p values of t-test for the analysed parameters in serum	111
Table D.2 : p values of t-test for the analysed parameters in heart	111
Table D.3 : p values of t-test for the analysed parameters in liver.....	112

LIST OF FIGURES

	<u>Page</u>
Figure 2.1 : Olive oil production	5
Figure 2.2 : Health effects of olive oil	8
Figure 2.3 : Main targets of oxidative stress	22
Figure 2.4 : Amino-thiol metabolism and GSH synthesis	27
Figure 4.1 : Mean ($\pm$ SEM) PON1 activity in serum of SAMR1 and SAMP8 mice fed a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either LP or HP .58	58
Figure 4.2 : Mean ($\pm$ SEM) vitamin C concentration (nmol/mg protein) in liver A) in heart B) of SAMR1 and SAMP8 mice	63
Figure 4.3 : Mean ($\pm$ SEM) vitamin E concentration (mmol/mg protein) in liver of SAMR1 and SAMP8 mice	64
Figure 4.4 : Relative mRNA expression of genes in vitamin E metabolism (normalized for GAPDH) of CYP3A (A), TTP (B), LPL (C), LRP1 (D), SRB1 (E), HMGCS (F), LDLR (G), ABCA1(H) in liver of SAMP8 and SAMR1 mice	66
Figure 4.5 : Relative mRNA expression (normalized for GAPDH) of Nrf2 (A), GST (B), γ -GCS (C), NQO- 1 (D), HO-1 (E), and PON-2 (F) in hearts of SAMP8 mice	68
Figure 4.6 : Protein levels of HO-1. Protein levels were determined via western blotting	69
Figure 4.7 : Relative mRNA expression (normalized for GAPDH) of Nrf2 (A), GST (B), γ -GCS (C), NQO- 1 (D), HO-1 (E), and PON-1 (F) in livers of SAMP8 and SAMR1 mice	70
Figure 4.8 : Protein levels of GST- α (A) and γ -GCS (B) in livers of SAMP8 and SAMR1 mice	71
Figure 4.9 : Relative mRNA expression (normalized for GAPDH) of SIRT-1 in livers and heart of SAMP8 and SAMR1 mice	72
Figure 4.10 : Relative mRNA expression (normalized for GAPDH) of SAP and CRP levels in livers of SAMP8 and SAMR1 mice	73
Figure A.1 : Representative chromatogram of the standards.....	104
Figure A.2 : Representative chromatogram of olive oil extract	104
Figure A.3 : Representative chromatogram of blank olive oil extract	105
Figure A.4 : Representative chromatogram of α - and γ -tocopherol	105
Figure B.1 : Calibration curve for FRAP assay	106
Figure B.2 : Calibration curve for TEAC assay	106
Figure B.3 : Calibration curve for the determination of total phenolics	106
Figure B.4 : Calibration curve for tyrosol	107
Figure B.5 : Calibration curve for hydroxytyrosol	107
Figure B.6 : Calibration curve for vanillic acid	107
Figure B.7 : Calibration curve for caffeic acid	107
Figure B.8 : Calibration curve for p-coumaric acid	108
Figure B.9 : Calibration curve for ferulic acid	108
Figure B.10 : Calibration curve for oleuropein	108

Figure B.11: Calibration curve for pinoresinol	108
---	-----

EFFECT OF A DIET RICH IN OLIVE OIL PHENOLICS ON AGE-RELATED CHANGES IN MOUSE MODEL OF AGEING AND HPLC ANALYSIS OF OLIVE OIL PHENOLICS WITH ELECTROCHEMICAL DETECTOR

SUMMARY

Olive oil is the primary source of fat in the Mediterranean diet, which has been associated with reduced risk of cardiovascular mortality, cancer incidence and mortality and incidence of Parkinson and Alzheimer diseases. Despite the scarcity of clinical studies investigating the underlying mechanisms, beneficial effects of the Mediterranean diet have partly been attributed to the use of olive oil. It is an important source of monounsaturated fatty acids (mainly oleic acid). However, it has been suggested that the beneficial effects of olive oil consumption in the prevention of chronic diseases may not only be attributed to its fatty acid composition but also to its high content of phenolics.

Tyrosol, hydroxytyrosol and oleuropein are the main hydrophilic phenolic compounds of olive oil, and their pharmacological activities reported in many *in vivo* and *in vitro* studies. They show beneficial effects either directly by their antioxidant properties or indirectly with their impact on cellular signaling pathways, expression of certain genes and acting as inhibitors or activators of regulatory enzymes. Epidemiological studies are performed in terms of their protective role on cardiovascular diseases, neurodegenerative diseases and cancer.

Olive oil phenolics are well known for its antioxidant activities. But their role in metabolic pathways, their effect on transcription factors, gene expressions need to be investigated. In this study it is aimed to study the effect of diet rich in olive oil phenolic compounds on age-related changes in heart and liver tissues of 9-10 weeks old female SAMP8 (Senescence Accelerated Mouse Prone 8) mice. We fed SAMP8 semisynthetic diets with 10% olive oil containing either high (HP) or low amounts of olive oil phenolics (LP) for 4.5 months. As a control SAMR1 (senescence-accelerated mouse-resistant 1) mice were used that have a normal ageing process. SAMP8 is a widely used human model of ageing due to their shortened lifespan and pathological disorders such as elevated lipid and protein oxidation, inflammation, mitochondrial dysfunction, early onset of atherogenesis. These properties render SAMP8 mice as a suitable rodent model for experimental ageing research.

Oxidative stress and inflammation play an important role in the pathogenesis of ageing and age-related disorders. Therefore in this study the parameters regarding oxidative stress and inflammation were investigated. As a biomarker of oxidative stress in both tissues, lipid oxidation and protein oxidation were determined by TBARS (Thiobarbituric acid reactive substances) assay and protein carbonyls, respectively. Also proteasomal activity of different subunits in the proteasomal system was measured. In serum of mice total cholesterol, total triglycerides, HDL cholesterol and paraoxonase-1 (PON-1) activity were determined. Nrf2-dependent gene expression may be impaired during the ageing process. In both tissues transcription factor Nrf2 and its target genes; glutathione-S-transferase (GST), γ -glutamyl-cysteinyl-synthetase (γ -GCS), NADPH quionone oxidoreductase (NQO1),

hemeoxygenase (HO-1), paraoxonase-1 (PON-1 in liver) and PON2 in heart were the main focus. Their amounts were determined both in mRNA level and protein level. Furthermore the level of antioxidant vitamins (vitamin C & E), genes that play a role in the metabolism of these vitamins and the parameters of redox status were determined.

As a result different results in each parameter were obtained depending on the response of the tissue to the oxidative stress. Generally higher level of oxidative stress biomarkers was detected in both tissues of SAMP8 mice as compared to SAMR1 mice. Proteasomal activity and redox status remained unchanged in both tissues and mice types as well.

Nrf2 as well GST, γ -GCS, NOQ1 and PON2 mRNA levels were significantly higher in heart tissue of the HP as compared to the LP group. The mRNA expression of HO-1 did not differ in heart with HP treatment. The mRNA expression levels of Nrf2 and its target genes did not differ between groups in liver. Only HO-1 mRNA and protein level decreased in SAMP8 LP group as compared to SAMR1 LP group but no effect of HP treatment was observed.

The mRNA expression level of inflammation related genes, SAP (Serum Amyloid P), CRP (C Reactive Protein) significantly increased and HO-1 significantly decreased in the liver of SAMP8 mice as compared to SAMR1 mice. Interestingly the level of two well known antioxidant vitamins (vitamin C & E) significantly increased in SAMP8 LP group. The expression of genes regarding ascorbic acid uptake and synthesis (sodium-dependent vitamin C transporter type 1-SVCT1, sodium-dependent vitamin C transporter type2-SVCT2 and L-gulonolactone oxidase-Gulo) were also analysed. The mRNA expression level of SVCT1 expression was significantly increased in SAMP8 mice but no change was determined in terms of SVCT2 and Gulo level.

Our data suggests that oxidative stress and inflammation may play an important role in the ageing of liver and heart in SAMP8 mice. Depending on the results, a diet rich in olive oil phenolics may prevent oxidative stress in the heart of SAMP8 mice by modulating Nrf2-dependent gene expression. As the two olive oil that were used in the feeding study compared, it has been observed that they have similar phenolic profile. On the other hand the HP oil contains approximately 40 fold higher hydroxytyrosol as compared to LP oil. Therefore the beneficial effects of the diet may be attributed to its hydroxytyrosol content. Vitamin C level may be a response of the body to oxidative stress it increased to protect the body homeostasis. This increase has reflected to vitamin E level, which was higher in SAMP8 LP mice.

Another aim of that study is to develop and validate a method with the sensitive electrochemical detector for the quantification of olive oil phenolics in oils. Electrochemical detectors provide a new analytical tool for resolving and accurately detecting trace amounts of any electroactive compound in a wide range of samples. In our validated method it is possible to analyze selected phenolic compounds (tyrosol, hydroxytyrosol, oleuropein, pinoresinol, caffeic acid, vanillic acid, p-coumaric acid, ferulic acid) in 17 minutes. This method was validated in terms of selectivity, lower limit of quantification and detection, accuracy, precision, recovery, short term stability, freeze thaw stability and post preparative stability. The phenolics were quantified in ng level. The recoveries of these phenolic compounds ranged between 72-96%.

The liver and heart are the major organs responsible for a number of physiological processes that are crucial to sustaining life. Most of the studies dealing with SAMP8 mice have focused on brain aging. Therefore our study could be one of the pioneer studies in liver and heart ageing and associated pathophysiological processes in SAMP8 mice. On the other hand in the literature there is only one study that uses

electrochemical detector in the quantification of olive oil phenolics but it is not validated. In this respect our validated method with good recovery of phenolics and lower limit of detection (ng level) will be beneficial for the future analysis of olive oil phenolics in shorter time.

ZEYTİNYAĞI FENOLİKLERİNCE ZENGİN DİYETİN YAŞLANDIRILMASI HIZLANDIRILMIŞ FARE MODELİNDE YAŞLANMAYA BAĞLI DEĞİŞİKLİKLER ÜZERİNE ETKİSİ VE ZEYTİNYAĞINDAKİ FENOLİK MADDELERİN ELEKTROKİMYASAL DEDEKTÖR İLE HPLC ANALİZİ

ÖZET

Zeytinyağı Akdeniz diyetinin temel yağ kaynağıdır. Zeytinyağı tüketimi kardiyovasküler hastalıklar, kanser, Parkinson ve Alzheimer gibi hastalıkların oluşum riskini ve bu hastalıklardan kaynaklanan ölüm riskini azaltmaktadır. Akdeniz diyetinin yararlı etkilerinin altında yatan mekanizmaları inceleyen klinik çalışmalar eksik olmasına rağmen, bu etkiler kısmen zeytinyağı tüketimine bağlanmaktadır. Zeytinyağı önemli bir doymamış yağ asidi (oleik asit) kaynağıdır. Ancak kronik hastalıkların önlenmesi ile ilişkili olarak zeytinyağı tüketiminin yararlı etkileri sadece yağ asidi bileşimine değil yüksek fenolik madde miktarına da bağlanmaktadır.

Tirozol, hidroksitirozol ve oleuropein zeytinyağında bulunan temel fenolik bileşenlerdir ve birçok in vivo ve in vitro çalışmada farmakolojik aktiviteleri belirtilmiştir. Bu bileşenler yararlı etkilerini direkt olarak antioksidan özellikleri ile veya indirekt olarak hücresel sinyal yollarını, bazı genlerin anlatımını etkileyerek, bazı düzenleyici enzimleri aktive ya da inhibe ederek göstermektedir. Bu bileşenlerin kardiyovasküler hastalıklar, nörodejeneratif hastalıklar ve kanser üzerine koruyucu etkileri ile ilgili olarak epidemiyolojik çalışmalar gerçekleştirilmektedir.

Zeytinyağı fenoliklerinin antioksidan aktivitesi çok iyi bilinmektedir. Ancak metabolik yollarındaki görevleri, transkripsiyon faktörleri ve gen anlatımı üzerine etkilerinin araştırılması gerekmektedir. Bu çalışmada fenolik maddelerce zengin zeytinyağı içeren diyetin 9-10 haftalık dişi SAMP8 (Senescence Accelerated Mouse Prone 8) farelerinin kalp ve karaciğer dokularında yaşlanmaya bağlı oluşan değişiklikler üzerine etkisinin incelenmesi amaçlanmıştır. SAMP8 fareleri 4.5 hafta boyunca düşük (DF) ya da yüksek fenolik (YF) madde miktarına sahip %10 zeytinyağı içeren yarı sentetik diyetlerle beslenmiştir. Kontrol olarak normal yaşlanma prosesine sahip SAMR1 (senescence-accelerated mouse-resistant 1) fareleri kullanılmıştır. SAMP8 fareleri düşük yaşam döngüsüne ve yüksek lipit ve protein oksidasyonu, iltihaplanma, mitokondri bozuklukları, erken aterogenez başlangıcı gibi patolojik bozukluklara sahip olduğundan dolayı insan yaşlanma modeli olarak sıklıkla kullanılmaktadır. Bu özellikler SAMP8 fare grubunu yaşlanma ile ilgili çalışmalarda kullanımını uygun kılmaktadır.

Oksidatif stres ve iltihaplanma yaşlanma ve yaşlanmaya bağlı bozuklukların patogeneğinde önemli rol oynamaktadır. Bu nedenle bu çalışmada oksidatif stres ve iltihaplanmaya bağlı parametreler incelenmiştir. Her iki dokuda da oksidatif stres biyomarkırı olarak lipit ve protein oksidasyonu sırasıyla TBARS ve protein karbonil metotları ile belirlenmiştir. Ayrıca proteazomal sistemdeki farklı altbirimlerin proteazomal aktivitesi ölçülmüştür. Farelerin serumunda toplam kolesterol, toplam trigliserit, HDL kolesterol ve paraoksonaz-1 aktivitesi belirlenmiştir. Yaşlanmaya bağlı olarak Nrf2 (nuclear factor erythroid 2-related factor 2) ye bağlı gen anlatımı bozulabildiğinden Nrf2 ve hedef genlerinin anlatımı; glutatyon-S-transferaz (GST), γ -glutamil-sisteinil sentetaz (γ -GCS), NADPH kuinon oksidoredüktaz (NQ1),

hemoksijenaz (HO-1), paraoksonaz-1 (PON-1 karaciğerde) ve kalpte PON2, hedef olarak seçilmiştir. Bu genlerin miktarları mRNA ve protein seviyesinde belirlenmiştir. Bunun dışında antioksidan vitamin miktarları (E ve C vitamini), bu vitaminlerin metabolizmasında rol oynayan genler ve redoks durumu parametreleri belirlenmiştir.

Sonuç olarak her parametrede dokuların oksidatif strese karşı yanıtına bağlı olarak farklı sonuçlar elde edilmiştir. Genel olarak SAMP8 farelerinin her iki dokusunda da oksidatif stres biyomarkırları SAMR1 farelerine göre yüksek saptanmıştır. Proteazomal aktivite ve redoks durumu parametreleri her iki doku ve fare tiplerinde değişmemiştir.

Nrf2 ve buna bağlı GST, γ -GCS, NOQ1 ve PON2 genlerinin mRNA seviyeleri kalpte YF grup farelerde DF grup farelere göre önemli derecede yüksek saptanmıştır. YF uygulaması ile HO-1 mRNA seviyesinde bir değişim gözlenmemiştir. Karaciğerde Nrf2 ve hedef genlerinin anlatımında gruplar arasında fark bulunmamıştır. Sadece HO-1 mRNA ve protein seviyesi SAMP8 DF grubunda SAMR1 DF grubuna oranla düşük saptanmıştır. Ancak YF uygulamasının bir etkisi saptanmamıştır.

Karaciğerde itihaplanma ile ilişkili olan Serum amyloid P (SAP) ve C Reaktif Protein (CRP) genlerinin mRNA anlatım miktarları SAMP8 DF farelerinde SAMR1 DF farelerine göre önemli derece artmış, HO-1 geninin mRNA miktarı ise azalmıştır. İlginç olarak iki iyi tanınan antioksidan vitaminlerin miktarı (E ve C vitamini) SAMP8 DF grubunda önemli derecede artmıştır. Sodyuma bağlı C vitamini taşıyıcısı-1 (SVCT1), sodyuma bağlı C vitamini taşıyıcısı-2 (SVCT2), L-gulonolakton oksidaz gibi askorbik asidin doku içine alımı ve sentezi ile ilişkili genlerin anlatımı analizlenmiştir. SVCT1 anlatım miktarı SAMP8 DF grubu farelerde önemli derece artmış ancak SVCT2 ve L-gulonolakton oksidaz açısından bir değişiklik görülmemiştir.

Elde edilen veriler SAMP8 farelerinin kalp ve karaciğer yaşlanmasında oksidatif stres ve iltihaplanmanın önemli rol oynayabileceğini göstermektedir. Sonuçlara bağlı olarak zeytinyağı fenolik maddelerince zengin diyetin SAMP8 farelerinde Nrf2 ve buna bağlı genlerin anlatımını düzenleyerek oksidatif stresi önleyebileceği söylenebilir. Besleme çalışmalarında kullanılan iki yağ örneği karşılaştırıldığında her ikisinin benzer profillere sahip olduğu görülmektedir. Diğer yandan YF yağ örneği DF yağ örneğine göre yaklaşık 40 kat daha fazla hidroksitirozol içermektedir. Bu nedenle YF içeren diyetin yararlı etkileri hidroksitirozol içeriğine de bağlanabilir. Karaciğerde SAMP8 farelerinde görülen yüksek C vitamini miktarı oksidatif strese karşı vücut dengesini sağlamak için vücudun bir cevabı olarak düşünülebilir. SAMP8 DF farelerinde görülen C vitaminindeki bu artış C vitamini tarafından indirgenen E vitamini miktarına da yansımıştır.

Bu çalışmanın diğer bir amacı da hassas elektrokimyasal dedektör ile zeytinyağındaki fenolik maddelerin belirlenmesi için metot geliştirmek ve metodun validasyonunu sağlamaktır. Elektrokimyasal dedektörler pekçok üründe iz miktardaki elektroaktif bileşiklerin doğru bir şekilde belirlenmesinde yeni bir analitik araç olarak görev yapmaktadır. Çalışmada valide edilmiş metot ile zeytinyağında bulunan seçilmiş fenolik maddelerin analizi (tirozol, hidroksitirozol, oleuropein, piniresinol, kafeik asit, vanilik asit, p-koumarik asit, ferulik asit) 17 dakika içinde yapılması mümkün olmuştur. Bu metot seçicilik, en düşük belirlenme limit değeri, geri kazanım, doğruluk, duyarlık, dondurma-çözündürme dayanıklılığı, kısa süreli dayanıklılık, örnekleyici içindeki dayanıklılık açısından valide edilmiştir. Fenolik maddeler ng seviyesinde belirlenmiştir. İncelenen fenolik maddelerin geri kazanımları %72-96 arasında saptanmıştır.

Kalp ve karaciğer hayatı sürdürmek için gerekli pekçok fizyolojik prosesten sorumlu organlardır. SAMP8 fareleri ile gerçekleştirilen pekçok çalışmada beyin yaşlanması üzerinde durulmuştur. Bu nedenle yapılan çalışma SAMP8 farelerinde kalp ve

karaciğer yaşılanması ve buna bağlı patofizyolojik prosesler açısından öncü bir çalışma olarak değerlendirilmektedir. Diğer yandan literatürde zeytinyağının fenolik maddelerinin elektrokimyasal dedektör ile belirlenmesi açısından sadece bir çalışma mevcuttur ancak metot valide edilmemiştir. Bu açıdan valide edilmiş, fenolik maddelerin geri kazanımı oldukça yüksek ve en düşük belirlenme limitinin ng seviyesinde olduğu metot gelecekte kısa sürede fenolik madde analizi açısından yararlı olacaktır.

1. INTRODUCTION

The 'Mediterranean Diet is rich in olive oil, fruits, vegetables, grains and legumes. A high degree of adherence to the traditional Mediterranean diet has been associated with a reduced risk of overall and cardiovascular mortality, cancer incidence and mortality, and incidence of Parkinson and Alzheimer disease (Trichopoulou et al., 2003; Sofi et al., 2008). Despite the scarcity of clinical studies investigating the underlying mechanisms, beneficial effects of the Mediterranean diet have partly been attributed to the use of olive oil (Alonso et al., 2004; Esposito et al., 2004; Estruch et al., 2006; Kontogianni et al., 2007). Olive oil is the primary source of fat in the Mediterranean diet.

Olive oil is obtained from the fruits of *Olea Europea L.* tree and it contributes ~4% of total vegetable oil production. More than 95% of world olive oil production is supplied by Mediterranean countries. Due to growing interest in Mediterranean diet and its health benefits, the production of olive oil is now expanding to non-traditional producers such as United States, Canada, Australia and Japan (Visioli et al., 2002).

Olive oil is composed of major and minor components. The major components represent more than 98% of the total oil weight and they include glycerols. High content of oleic acid is unique to olive oil. Compared to polyunsaturated fatty acids, oleic acid is monounsaturated, making it much less susceptible to oxidation and contributing to the antioxidant action, high stability and long shelf life of olive oil (Owen et al., 2000). The non-glycerol or unsaponifiable fraction consists of 0.4-2% including more than 230 chemical compounds such as sterols, aliphatic alcohols, triterpene alcohols, hydrocarbons, pigments, volatile compounds, and antioxidant compounds (Servili et al., 2004; Tripoli et al., 2005).

Olive oil is known to have antioxidant (Liu et al., 2007; Loru et al., 2009), anti-inflammatory (Bogani et al., 2007; Zhang et al., 2009), anti-atherogenic (Visioli and Galli, 2000), anti-ulcer (Serrano et al., 1997), anti-cancer (Della Regione et al., 2000; Quiles et al., 2002; Gill et al., 2005), anti-microbial (Medina et al., 2006), diabetic (Soriguer et al., 2004; Al-Azzawie et al., 2006; Rigacci et al., 2009), neuroprotective (Schaffer et al., 2007; Young et al., 2008), skin protective (Perugini et al., 2008) and anti-ageing (Hao et al., 2009) effects. The overall beneficial effects

of virgin olive oil have been linked to both oleic acid and its antioxidant components. The main antioxidants of olive oil are β -carotenes and phenolic compounds including lipophilic and hydrophilic phenols (Boskou, 1996). While the lipophilic phenols, among which tocopherols can be found in other vegetable oils, some hydrophilic phenols of olive oil (including phenolic acids, secoiridoid compounds and derivatives) are not generally present in other oils and fats. Moreover, the hydrophilic phenols of olive oil constitute a group of secondary plant metabolites which show sensory and health properties (Servili et al., 2004).

Tyrosol, hydroxytyrosol and oleuropein are the main phenolic compounds of olive oil, and their pharmacological activities reported in many *in vivo* and *in vitro* studies. They show beneficial effects either directly by their antioxidant properties or indirectly with their impact on cellular signaling pathways, expression of certain genes and acting as inhibitors or activators of regulatory enzymes. Epidemiological studies are performed in terms of their protective role on cardiovascular diseases, neurodegenerative diseases and cancer.

The susceptibility to these diseases increases with ageing. Aging is associated with several changes in structure and function of different organs and tissues and it is characterized as a progressive decline in biological functions with time (Farooqui and Farooqui, 2010). One of the most accepted theories of aging is the free radical theory, which postulates that aging is the result of the accumulation of cellular and tissue damage caused by the continuous and endogenous generation of reactive oxygen species (ROS) (Sohal, 2002; Harman, 2003; Barja, 2004). As the amount of ROS increases oxidative stress occurs, that contributes to the pathogenesis and progression of age related disorders (Kregel and Zhang, 2007).

Another theory for ageing is the molecular inflammation hypothesis suggesting that there is link between chronic, low grade inflammation and the pathogenesis of age-related diseases, it is based on dysregulation of the immune system with age and altered redox status during aging. Both processes lead to activation of a wide variety of inflammatory mediators through overproduced reactive species during aging (Chung et al., 2009).

There is a great interest in the development of anti-ageing therapies, protecting from the deleterious effects of ageing, and phytochemicals, dietary interventions containing antioxidants have been widely investigated in this respect. Olive oil, a food naturally containing high amounts of phenolic compounds, may be an alternative. To understand the effect of olive oil phenolics on ageing *in vivo* studies

are warranted. Researchers have demonstrated that several genes may modulate aging and influence longevity. Olive oil phenolic compounds are well known for their antioxidant activities. But their role in metabolic pathways, their effect on transcription factors, gene expressions needs to be investigated.

In our study it is aimed to study the effect of diet rich in olive oil phenolic compounds on age-related changes in heart and liver tissues of SAMP8 mice, a widely used rodent model for experimental ageing research, which have a shorter life span and pathological disorders. As oxidative stress and inflammation play an important role in the pathogenesis of ageing and age-related disorders, the parameters regarding these factors were investigated. In both tissues Nrf2 and the genes that are controlled under that transcription factor was the main focus. Furthermore the level of antioxidant vitamins (vitamin C and vitamin E), biomarkers of redox status were measured. As a biomarker of oxidative stress, lipid oxidation was determined by TBARS assay, protein oxidation was determined by measuring protein carbonyl levels. Also proteasomal activity of different subunits in the proteasomal system was measured.

Another aim of that study is to develop and validate a method with the sensitive electrochemical detector for the quantification of phenolics in olive oil.

2. LITERATURE

2.1 Production of Olive Oil

The most vegetable oils are extracted from seeds by solvents. On the other hand olive oil is produced by a cold-press mechanical process that preserves both the chemical nature of the oil and the natural antioxidants that are produced in response to environmental stress which is suggested as the uniqueness of olive oil (Visioli and Galli, 2000).

The first step in olive oil production is washing of the olives in order to remove any foreign material that could damage machinery or contaminate the oil. Then the olives are crushed to produce a paste with easily extracted oil droplets. A step called malaxation takes place, where olive paste is mixed and prepared for separation of the oil from the pomace. The next step is separating the oil from the paste and fruit water (water of vegetation). The oil can be extracted by pressing, centrifugation, percolation, or through combinations of the different methods. In the last step oil is separated from the water. The figure 2.1 shows olive oil production with classical, two phase and three phase systems.




Figure 2.1: Olive oil production (Boskou, 2006).

2.2 Chemical Composition of Olive Oil

Olive oil is composed of major and minor components. The major components include glycerols and represent more than 98% of the total oil weight. The non-glycerol or unsaponifiable fraction consists of 0.4-2% (Servili et al., 2004; Tripoli et al, 2005). The chemical composition of olive oil is given in Table 2.1.

Table 2.1: Chemical composition of olive oil.

Saponifiable fraction (98-99%)	Unsaponifiable fraction (~2%)
Main fatty acids present in triacylglycerols Miristic acid (14:0) Palmitic acid (16:0) Palmitoleic acid (16:1) Stearic acid (18:0) Oleic acid (18:1) Linoleic acid (18:2) Linolenic acid (18:3)	Non glyceride esters (alcoholic and sterol compounds, waxes) Aliphatic alcohols Triterpene alcohols Sterols (β -sitosterol, campesterol, stigmasterol) Hydrocarbons (squalene, lycopene, β -carotene) Pigments (β -carotene, chlorophylls) Lipophilic phenolic (Vitamin E) Hydrophilic phenolics (phenolic acids, phenolic alcohols, secoiridoids, lignans, flavones) Volatile compounds

Olive oil is graded in six categories: extra virgin olive oil (EVOO), virgin olive oil (VOO), refined olive oil, olive oil, refined residue oil, and olive residue oil. These oils differ in their acidity, way of production and micronutrients content. Among them VOO and EVOO are attributed to health effects of olive oil.

2.2.1 Major compounds

Triacylglycerols are constituted by most of the 12 identified fatty acids (C14-C24), although only 6 are major compounds. These are; oleic acid (55.23-86.64%), palmitic acid (6.30-20.93%), linoleic acid (2.7-20.24%), stearic acid (0.32-5.33%), palmitoleic (0.32-3.52%), and linolenic acid (0.11-1.52%) (Garcia-Gonzales et al., 2008). Virgin olive oil is unique with respect to the high oleic acid content because the majority of seed oils are composed primarily of polyunsaturated fatty acids, including the essential omega-6 fatty acid, linoleic acid. Compared to polyunsaturated fatty acids, oleic acid is monounsaturated, making it much less susceptible to oxidation and contributing to the antioxidant action, high stability and long shelf life of olive oil (Owen et al., 2000).

2.2.2 Minor compounds

Minor compounds of virgin olive oil are present in low amount and include more than 230 chemical compounds such as sterols, aliphatic alcohols, triterpene alcohols,

hydrocarbons, pigments, volatile compounds, lipophilic and hydrophilic phenolic compounds (Servili et al, 2004).

2.2.2.1 Lipophilic phenolics (Vitamin E)

Vitamin E is the most abundant lipophilic phenolic. It is a general name of four tocopherol (namely α -, β -, γ -, and δ -) and four tocotrienol isoforms (namely α -, β -, γ -, and δ -). Virgin olive oil contains high amount of α -tocopherol in range of 50-250 mg/kg followed by β - and γ -tocopherols (García-González et al., 2008).

2.2.2.2 Sterols

β -sitosterol is represent the sterol content of virgin olive oil with 90-95% that is followed by campesterol and stigmasterol with 3% and 1% respectively (Gutierrez et al., 1999). The sterol amount of virgin olive oil is approximately 1600 mg/kg (Covas et al., 2006).

2.2.2.3 Hydrocarbons

Squalene is the most important hydrocarbon of virgin olive oil which constitutes around 50% of the unsaponifiable matter (García-González et al., 2008). It is the metabolic precursor of cholesterol and other sterols and it has a role in the synthesis of steroid hormones. It is found in olive oil (~0.7%) in higher amounts as compared to other vegetable oils (0.002-0.03%) (Reedy and Couvreur, 2009).

2.2.2.4 Pigments

Chlorophyll and carotenoid pigments are responsible for the color of virgin olive oil, ranging from yellow-green to greenish gold. The chlorophyll pigments are responsible for the greenish hues of virgin olive oil, with pheophytin a being a major compound in this regard, while the major “yellow” pigments are lutein and β -carotene. Carotenoids are pigments whose content does not usually exceed 10 mg/kg, with lutein being the major carotenoid followed by β -carotene and other xanthophylls. Xanthophylls esterified with fatty acids play an important role in the stability of carotenoids, stabilizing the oil colors and ensuring a sufficient level of substances with antioxidant activity (García-González et al., 2008).

2.2.2.5 Volatile compounds

Volatile are responsible for the olive oil flavor and aroma. They are the main compounds responsible for the “green” and fruity desirable aroma attributes of olive oil. On the contrary, the low-quality olive oil have complex profiles composed of a

large number of volatiles responsible for off flavors such as rancid, mustiness, fusty and muddy sediment (García-González et al., 2008).

2.2.2.6 Triterpenes

Triterpenoic alcohols (erithrodiol and uvaol) and triterpenoic acids (maslinic, oleanolic, ursolic and betulinic acids) are the most relevant triterpene molecules in virgin olive oil. They are concentrated in the pericarp of olives. Therefore it is found in virgin olive oil in less amount (Covas et al., 2006).

2.3 Health Benefits of Olive Oil

Olive oil is characterized with its high content of monounsaturated fats and is also a good source of phytochemicals including phenolic compounds, squalene and vitamin E. It is also considered as a “natural functional food” or “liquid gold” with protective effects on oxidative stress and age-related disorders such as cancer, cardiovascular diseases, ulcer, diabetes, alzheimer and dementia. The health effects of olive oil is given in Figure 2.2. Many intervention studies (Tzonou et al., 1993; Braga et al., 1998; Stoneham et al., 2000; Fung et al., 2009; Perona et al., 2010; Martinez-Gonzalez et al., 2011) demonstrate that the lower incidence of cancer and cardiovascular diseases is attributed to high consumption of olive oil in Mediterranean countries.




Figure 2.2: Health effects of olive oil.

The overall beneficial effects of virgin olive oil have been linked to both its monounsaturated fatty acids (oleic acid) and its antioxidant components. It cannot

be explained by the effect of a single component. These compounds show their effect by different mechanism of actions as summarized in Table 2.2.

In recent years many epidemiological and clinical studies have been conducted with phenolic compounds of virgin olive oil and more emphasis will be given to these compounds in the next sections including their health benefits, technological role in olive oil and their analysis techniques.

Table 2.2: Mechanism of actions of olive oil components in their health effects.

Compound	Mechanism of action	Reference
Sterols	Protect against membrane lipid peroxidation, anti-inflammatory effect, improvement of endothelial dysfunction, decreases plaque formation, decreases LDL cholesterol, total cholesterol and triglycerides, anti-carcinogenic effect against lung, breast, stomach cancers via decreasing the growth of tumors, increasing the immune response and antioxidant enzyme activity	Mendilaharsu et al., 1998; De Stefani et al., 2000; van Rensburg et al., 2000; Devaraj et al., 2006; Raitakari et al., 2008; Hallikainen et al., 2006; Seppo et al., 2007
Skualen	Protection against UV radiation and radioactivity	Reedy and Couvreur, 2009
Carotenoids	Protective effect against cancer and cardiovascular diseases, antioxidant effect	Bernstein et al., 2002; Voutilainen et al., 2006
Luteolin	Protective against cataract and macular degredation, antioxidant and anticarcinogenic effect, protection of DNA from oxidation	Trumbo and Ellwood 2006; Santocono et al., 2007; Kim et al., 2007
Chlorophylls	Antioxidant effect, protective against cancer	Kamat et al., 2000; Lanfer-Marquez et al., 2005; Ferruzzi and Blakeslee, 2007
α -Tocopherol	Antioxidant effect, prevent oxidation of lipoproteins and biological membranes, prevent atherosclerosis, reduce breast cancer risk, protection of DNA from oxidation, neuroprotection	Rimm et al., 1993; Saldeen et al., 1999; Kline et al., 2003
Oleic acid	Decreases triacylglycerol and cholesterol, anti-inflammatory and antihypertensive effects, protects cell membrane and lipoproteins from oxidative stress, lowers the risk of colon, lung, breast, coleractal, and prostate cancers, improve insulin sensitivity, lowers plasma glucose and insulin levels	Yamaki et al., 2002; Lopez et al., 2008; Paniagua et al., 2007; Barberger-Gateau et al., 2007; Solfrizzi et al., 2006
Triterpenes	Antioxidant, anti-inflammatory, vasodilatory, cardiotonic effects	Somova et al., 2003; Martinez-Gonzalez et al., 2008

2.4 Hydrophilic Phenolic Compounds

Virgin olive oil contains different classes of phenolic compounds such as phenolic acids, phenolic alcohols, flavonoids, hydroxy-isocromans, secoiridoids, and lignans as reported in Table 2.3. The most abundant phenols are the secoiridoids. These compounds, e.g., oleuropein and ligstroside, are derivatives of the secoiridoid glucosides of olive fruits. Breakdown products of two major phenolic constituents of the olive fruit, oleuropein and ligstroside, form the majority of the phenolic fraction. There are at least thirty-six structurally distinct virgin olive oil phenolics that have been identified (Cicerale et al., 2009). They can be grouped according to their similar chemical structure in the following groups:

Phenolic acids with the basic chemical structure of C6-C1 (benzoic acids) and C6-C3 (cinnamic acids), such as caffeic, vanillic, p-coumaric, o-coumaric, protocatechuic, sinapic, and p-hydroxybenzoic acid, are the first group of phenols observed in olive oil.

Phenolic alcohols are the compounds with hydroxyl group attached to an aromatic hydrocarbon group. Hydroxytyrosol and tyrosol are the main phenolic alcohols in virgin olive oil. Secoiridoids are characterized by the presence of either elenoic acid or elenoic acid derivatives in their molecular structure (Carrasco-Pancorbo et al., 2005). Hydroxyl-isocromans are 3,4-dihydroxy-1H-benzopyran derivatives (Bianco et al., 2001). Flavonoids are compounds containing two benzene rings joined by a linear three carbon chains, lignans are compounds which structure based on the condensation of aromatic aldehydes (Carrasco- Pancorbo et al., 2005). They are the second prevalent group in virgin olive oil.

Table 2.3: Phenolic compounds in virgin olive oil (Carrosco-Pancorbo et al., 2005; Servili et al., 2004).

Phenolic acids and derivatives
Gentisic acid Vanillic acid p-coumaric acid o-coumaric acid Gallic acid Caffeic acid Protocatechuic acid Ferulic acid Cinnamic acid Benzoic acid p-hydroxybenzoic acid Sinapinic acid 4-Acetoxyethyl-1,2-dihydroxybenzene 2-(4-hydroxyphenyl)ethyl acetate
Phenolic alcohols
(3,4-dihydroxyphenyl)ethanol (3,4 DHPEA) (p-hydroxyphenyl)ethanol (p-HPEA) (3,4-dihydroxyphenyl)ethanol glucoside
Secoiridoids
Dialdehydic form of decarboxymethyl elenoic acid linked to 3,4-DHPEA (3,4 DHPEA-EDA) Dialdehydic form of decarboxymethyl elenoic acid linked to p-HPEA (p-HPEA-EDA) Oleuropein aglycone (3,4 DHPEA-EA) Oleuropein p-HPEA derivative Dialdehydic form of oleuropein aglycone Dialdehydic form of ligstroside aglycone Ligstroside aglycone Oleochanthal
Lignans
(+)-1-Acetoxypinoresinol (+)-Pinoresinol (+)-1-Hydroxypinoresinol
Flavonoids
Apigenin Luteolin (+)-Taxifolin
Hydroxyisochromans
1-phenyl-6,7-dihydroxy-isochroman 1-(3-methoxy-4-hydroxy)phenyl-6,7-dihydroxy-isochroman

2.5 Factors that Affect the Phenolic Content of Olive Oil

The qualitative and quantitative composition of virgin olive oil hydrophilic phenols is strongly affected by the agronomic and technological conditions of production.

2.5.1 Agronomic conditions

Among agronomic parameters, the most studied aspects include cultivar, geographical origin, fruit ripening, climatic conditions of production and some agronomic techniques such as the irrigation.

As reported, phenolic composition of fruit is affected by the cultivar (Briante et al., 2002; Gomes-Alonso et al., 2002; Pinelli et al., 2003). Each olive cultivar has different phenolic content (Esti et al., 1998). For example Italian “coratina” cultivar is known to have higher phenolic content (Aparicio and Luna, 2002).

Geographical origin has a great influence on phenolic composition of virgin olive oils (Criado et al., 2004). Depending on the olive grove, climate and altitude change, thus climatic conditions (rainfall, temperature, humidity) differ.

In some aspects controversy results are obtained as there are many other factors that may affect the results. Several studies related to the relationships between irrigation during olive growing and phenolic concentration of olive oil. For instance Motilva et al. (1999) concluded that hydrophilic phenols in oil increase when the olives were grown under conditions of deficit of irrigation whereas Ismail et al. (1999) reported that the highest level of hydrophilic phenols in oil was obtained from regularly irrigated olives. Using a linear irrigation strategies Tovar et al. (2001), found a negative correlation between the level of water used and the concentration of secoiridoid derivatives of virgin olive oil, on the contrary the concentration of lignans was lower in the oil from the least irrigated treatment.

Olive ripening stage is another factor for the phenolic content. As the maturity increases, the amount of total phenols decreases in almost all samples (Beltran et al., 2005; Ayton et al., 2006). During maturation, the esterase activity causes degradation of oleuropein, the main phenolic compound present in unripened olive fruit. Therefore, it is important to establish the optimal harvest time, which should correspond to the optimal phenol content and the best quality of oils (Skevin et al., 2003).

2.5.2 Technological Conditions

Since the occurrence of hydrophilic phenols in virgin olive oil is strictly related to the activities of various endogenous enzymes of olive fruit, their concentration in the oil is strongly affected by the extraction conditions. Crushing and malaxation are the most important critical points of the oil mechanical extraction process (Servili et al, 2004). The amount of phenolics decreases as the time and the temperature of malaxation increases (Servili et al., 2003). Secoiridoid aglycons are originated during crushing from the hydrolysis of their glucosides, oleuropein, demethyloleuropein and ligstroside with the activity of endogenous glucosidases. It is reported that olives which were blanched before crushing to inactivate endogenous glycosidases the concentration of oleuropein and demethyloleuropein was not significantly modified (Servili et al., 2002).

Extraction systems, such as pressure and centrifugation, play an important role in the oil phenolic composition. Oil obtained by pressure or two phase centrifugation systems that do not require addition of water, shows higher phenolic concentration in comparison to the one obtained by the traditional three phase centrifugation system (Kaula et al., 2006). In general, the concentration of phenols in virgin olive oils may range between 40 and 900 mg/kg or up to 1000 mg/kg (Montedoro et al., 1992).

Apart from their nutritional value phenolic compounds play an important role in stability, and sensory properties of olive oil.

The phenolic constituents confer a bitter, astringent, throat-catching and pungent taste to the oil. Oils produced from greener olives, which contain high amount of phenolic compounds, have “fruity” and complex aroma. Conversely, “sweet” oils are almost devoid of phenols. On the other hand very high load of phenols may result in an excessive and unpleasant bitterness and is not synonymous with quality (Visioli and Galli, 2000). Oleuropein is the bitter principle of olives and is found in virgin olive oil as such and in its aglycone form. It undergoes hydrolysis and yields several simpler molecules (simple phenols) that build up the well-known olive oil complex taste. These modifications enhance the aroma and flavor of olive oils. Also strong correlation is reported between the bitter and pungent sensory notes and the ligstroside derivatives such as *p*-HPEA-EDA and 3,4-DHPEA-EDA (Gutiérrez-Rosales et al, 2001; Tovar et al., 2001).

Phenolic compounds also affect the stability of virgin olive oil. During storage, fatty acids in oils undergo oxidative degradation and odorless molecules, such as

hydroxyperoxides and secondary products, are produced. It is stated that virgin olive oil is highly stable to oxidation and has a longer shelf life as compared to other edible vegetable oils, due to its fatty acid composition, mainly high monounsaturated fatty acid content, and to the presence of minor compounds (lipophilic tocopherols and hydrophilic phenolic compounds) that have a major role in preventing oxidation. Particularly phenolic compounds contribute significantly to its prolonged shelf life and among them the secoiridoid derivatives show a large capacity to protect virgin olive oil from oxidative degradation (Gutierrez et al., 2001; Mateos et al., 2003; Bendini et al., 2006). Phenolic compounds can inhibit oxidation by a variety of mechanisms based on radical scavenging, hydrogen atom transfer, and metal-chelating attributes (Huang et al., 2005; Prior et al., 2005). Beside it is reported that virgin olive oil phenolics contribute to stability of olive oil under deep-frying conditions (Quiles et al., 2002; Napolitano et al., 2008; Casal et al., 2010).

2.6 Health Effects of Olive Oil Phenolics

Tyrosol, hydroxytyrosol and oleuropein are the main phenolic compounds of virgin olive oil, and their pharmacological activities reported in many studies.

Hydroxytyrosol is the principal phenolic compound found in olive oil and is known to possess strong antioxidant scavenging abilities. It has been reported that hydroxytyrosol inhibits low density lipoprotein (LDL) oxidation (Visioli, et al., 1998a), platelet aggregation (Petroni et al., 1995) and endothelial cell activation (Carluccio et al., 2003), accumulation of thromboxane and production of proinflammatory leukotrienes, cytokines as well as scavenge superoxide anion and hypochlorous acid (Visioli et al., 2002; Zhang et al., 2009). Hydroxytyrosol possesses a marked antioxidant activity, scavenging peroxy radicals (Saija et al., 1998), other ROS (Gordon et al., 2001) and RNS (Deiana et al., 1999; de la Puerta et al., 2001), or breaking peroxidative chain reactions and preventing metal ion catalyzed production of reactive oxygen species (Gutierrez et al., 2001). Hydroxytyrosol is highly protective against the peroxynitrite-dependent nitration of tyrosine and DNA damage by peroxynitrite in vitro (Deiana et al., 1999).

Oleuropein has been shown to possess several pharmacological properties, including antioxidant (Visioli et al., 2002), anti-inflammatory (Visioli et al., 1998a), anti-atherogenic (Carluccio et al., 2003) and anti-cancer (Owen et al., 2000) activities. In addition, oleuropein has been shown to be cardioprotective against acute adriamycin cardiotoxicity (Andreaddou et al., 2006) and to exhibit anti-ischemic and hypolipidemic activities (Andreaddou et al., 2006). It has been suggested to

reduce atherosclerosis development, improve antioxidant status (González-Santiago et al., 2006), increase iNOS activity by scavenging superoxide anion (Visioli et al., 1998b), increase resistance of LDL to oxidation (Coni et al., 2000). Both oleuropein and hydroxytyrosol have been proven to be more potent antioxidants than vitamin E and BHT (Visioli et al., 1998b).

Tyrosol is the most abundant biophenol in olive oil. Tyrosol is reported to scavenge peroxynitrite (de la Puerta et al., 2001) and O^{2-} (Bertelli et al., 2002). In addition to its antioxidative effects, it has been reported that tyrosol inhibits lipopolysaccharide (LPS)-induced cytokine release from human monocytes (Giovannini et al., 2002) and LPS-induced leukotriene B4 release in human mononuclear cells (de la Puerta et al., 1999). Although it has a weaker antioxidant capacity, it has been reported that tyrosol can restore antioxidant defense, modulate ROS and NO production, COX and lipoxygenase pathways (Moreno, 2003; di Benedetto et al., 2007). Tyrosol is able to inhibit LDL-cholesterol oxidation and to prevent the modification of the apoproteic moiety (Caruso et al., 1999). Tyrosol has also been effective in inhibiting leukocyte 5-lipoxygenase (de la Puerta et al., 1999), protecting the CaCo-2 intestinal mucosa cells against the cytostatic effects produced by oxidized LDL (Giovannini et al., 1999) and inhibiting lipopolysaccharide (LPS)-induced cytokine release from human monocytes (Giovannini et al., 2002). Moreover, it showed a neuroprotective effect against the peroxynitrite-induced neuronal oxidative damage (Vauzour et al., 2007) and in the focal cerebral ischemia/reperfusion rat model (Bu et al., 2007). Some other pharmacological effects of these main phenolics are summarized in Table 2.4.

Table 2.4: Health effect of the main olive oil phenolic compounds.

Phenolic compound	Mechanism of action	Protective effect	Conditions (Cell/mouse/human)	Level (in vivo/ invitro)	Reference
Hydroxytyrosol	Antioxidant	Scavenge hydrogen peroxide	Human neutrophils	In vitro	O'Dowd et al., 2004
	Antioxidant	Protect from acrolein-induced oxidative stress and mitochondrial dysfunction	Retinal pigment epithelial cells	In vitro	Liu et al., 2007
	Antioxidant, antiageing	Induce Phase II enzymes, enhance mitochondrial biogenesis and improve mitochondrial function, strengthen the cell's antioxidant defenses	3T3-L1 adipocytes	In vitro	Hao et al., 2009
	Antioxidant	protect cells from H ₂ O ₂ -induced damage via inhibiting production of MDA, hydroperoxides	Renal proximal tubular cells	In vitro	Loru et al., 2009
	Antidiabetic, antioxidant	inhibit hyperglycemia and oxidative stress induced by diabetes, increase antioxidant enzyme activities, decreases lipid peroxidation	Diabetes rat model	In vivo	Hamden et al., 2009
	Anticancer	lower the levels of hydroxyperoxides, DNA damage and mRNA levels of classic glutathione peroxidase	Human prostate cells	In vitro	Quiles et al., 2002
	Anticancer	antiproliferative activity, induce apoptosis	White blood cell from promyelocytic leukemia	In vitro	Della Regione et al., 2000
	Antioxidant, skin protective	Protect against UV-B induced DNA damage	HaCaT cells	In vitro	Guo et al., 2010
	Anti-inflammatory	Inhibit pro-inflammatory cytokines, iNOS and COX-2 expression	Human monocytic cells	In vitro	Zhang et al., 2009
	Antioxidant, Neuroprotective	Protect against DNA damage, quench radicals	Murine-dissociated brain cells	In vitro	Young et al., 2008

Table 2.4 (continued) : Health effect of the main olive oil phenolic compounds.

Phenolic compound	Mechanism of action	Protective effect	Conditions (Cell/mouse/human)	Level (in vivo/ invitro)	Reference
Hydroxytyrosol	Antioxidant, Neuroprotective	Attenuate Fe ²⁺ - and nitric oxide (NO)-induced cytotoxicity, prevents oxidative stress-induced loss of cellular ATP levels, decreases basal and iron-stimulated lipid peroxidation	PC-12 cells and mouse	In vitro and ex-vivo	Schaffer et al., 2007
Oleuropein	Antidiabetic, antioxidant	Decrease lipid peroxidation, inhibiting hyperglycemia , increase antioxidant enzyme activities	Diabetic rabbits	In vivo	Al-Azzawie et al., 2006
	Skin protective	Protect against UV-induced erythema	20-30 yrs females	In vivo	Perugini et al., 2008
	Neuroprotective	Stabilize amyloid- β structure and prevent the formation to a toxic β -sheet conformation	Model system	In vitro	Bazoti et al., 2008
	Antidiabetic	Interfere with human islet amyloid polypeptide (hIAPP) and protect against aggregate cytotoxicity in type II diabetes	Rat RIN-5F insulinoma pancreatic β cells	In vitro	Rigacci et al., 2009
Tyrosol	Neuroprotective	Protect against the sensory motor dysfunction, neuroprotection	Stroke rat model	In vivo	Bu et al., 2007
Oleocanthal	Anti-inflammatory	Inhibit COX-1 and COX-2	Model system	In vitro	Smith et al., 2005
	Neuroprotective	Alter aggregation of amyloid- β protein via changing its structure, immunoreactivity	Model system	In vitro	Pitt et al, 2009
		Inhibit Tau fibrilization	Model system	In vitro	Li et al., 2009
Pinoresinol rich olive oil	Anticanceronic	Decrease cell viability, induce apoptosis	Colon cancer cell line	In vitro	Fini et al., 2008

2.7 Analytical Determination of Phenolic Compounds In Olive Oil

The identification and quantification of phenolic compounds are of great interest due to its nutritional and technological significance. The first step is the extraction of phenolic compounds. The most prevalent methods are solid phase extraction (SPE) and liquid-liquid extraction (LLE). Different cartridges are used in SPE including C8, C18, NH₂, and diol. But more promising results are obtained with diol cartridges being the most suitable cartridge for phenolic analysis. In terms of recovery comparable results are obtained in the literature. But SPE is more advantageous than LLE in terms of cleanliness, simplicity, time of analysis, less amounts of solvents and sample required.

For the quantitative determination of total phenols the most widely used method is Folin-Ciocalteu Method which is based on the reaction Folin-Ciocalteu reagent with the hydroxyl groups of phenolic compounds. After the colour reaction the absorbance is measured spectrophotometrically (Singleton et al., 1999). It is a simple and fast method but has low specificity as the colour reaction can occur with any oxidizable phenolic hydroxyl group.

Apart from that various methods have been developed to analyze the qualitative and quantitative occurrence of phenolic compounds. The simple techniques include Thin Layer Chromatography (TLC) (Ragazzi et al., 1973), reversed phase High Performance Liquid Chromatography (HPLC) coupled with Ultraviolet (UV) (Tsimidou et al., 1992), Diode Array Detector (DAD) (Pirisi et al., 1997; Servili et al., 1999; Esti et al., 1998), ECD (Electrochemical Detector) (Brenes et al., 1999), Gas Chromatography (GC) alone (Solinas et al., 1982) or coupled with Mass Spectrometry (GC-MS) (Liberatore et al., 2001; Marsilio et al., 2001) and Capillary Electrophoresis (CE) (Bendini et al., 2003; Bonoli et al., 2004). Beside phenolic compounds can be determined by NMR (nuclear magnetic resonance) (Angerosa et al., 1995; Montedoro et al., 1993, Christophoridou et al., 2005) and IR (Infrared) (Montedoro et al., 1993) techniques.

Among them LC-MS determination of phenolic compounds is the most prevalent technique which can determine the compounds in ng range. However it is hard to interpret MS results, without having any additional information about the compound.

Another technique that is comparable with LC-MS in terms of sensitivity is coulometric electrode array detector (a type of Electrochemical Detector) coupled to HPLC. Because of a high selectivity and sensitivity, HPLC with electrochemical

detection has become increasingly popular for the analysis of natural antioxidants in food and beverages. It is frequently used in the analysis of phenolic compound in fruits (Aaby et al., 2005; Novak et al., 2008), vegetables (Xu et al., 2006), wort (Vanbeneden et al., 2006), beer (Řehov et al., 2004), beverages (Peng et al., 2005; Wu et al., 2007; Novak et al., 2010) and pharmaceutical analyses. Electrochemical detectors provide a new analytical tool for resolving and accurately detecting trace amounts of any electroactive compound in a wide range of samples. However in the literature there is only one study that uses that detector in quantification of olive oil phenolics.

2.8 Principle of the Electrochemical Detector

EC detectors apply a voltage at an electrode surface over which the HPLC eluent flows. Electroactive compounds eluting from the column either oxidize or reduce generating a current peak in real time. The amount of current generated depends on both the concentration of the analyte and the voltage applied, with each compound having a specific voltage at which it begins to oxidize (Svendsen 1993).

There are two types of electrochemical detectors that can be coupled to LC. These are amperometric and coulometric detectors. Both are based in the same principle of oxidation/ reduction of the analyte induced by an electrode held at a fixed potential. In an amperometric detector, mobile phase containing analytes flows over a planar electrode and only partial oxidation/reduction occurs, whereas in a coulometric detector the carrier passes through a porous electrode, resulting in an increased reactive surface and therefore almost complete oxidation/ reduction. Thereby the sensitivity of the detector increases (Penalvo and Nurmi, 2005).

The coulometric array detector simply consists of a series of coulometric electrode pairs placed in series performing a multichannel (or array) detection. The system is based on analytical cells containing a reference electrode that sets the electrochemical zero and four working electrodes that measure the redox reaction of interest. Each analytical cell therefore provides four channels and the systems available at present can bear up to four cells. It can be built up to 16 channels (Penalvo and Nurmi, 2005).

2.9 Advantages of Electrochemical Detector

-Enhanced Resolving Power: In HPLC with UV or DAD some contaminants in the sample co-elute with peaks of interest even after optimizing the chromatographic

conditions. Under, these conditions coulometric EC array detectors provide the separation and quantification of the co-eluting compounds have different oxidation potentials (Achilli et al., 1993).

-Simplified Sample Preparation: With the increased resolving power of EC array detectors it is possible to separate interfering compounds from the compound of interest. Hence, in many instances sample clean-up procedures can be simplified (Svendsen, 1993).

-The HPLC with ED is more sensitive, selective, and has a lower detection limit than that obtained by diode array detection.

-Confirmation of Compound Purity: When compounds pass through the EC array they are normally detected on three contiguous electrodes. The first electrode will oxidize a small portion of the compound, the second or dominant electrode oxidizes a large portion of the compound and the third electrode oxidizes the remainder of the compound. Pure standards eluting at a given retention time will give a predictable response at all three electrodes and the ratio across these three electrodes remains constant and is independent of concentration. In real samples, compounds eluting at a retention time matching that of the standard can also be 'ratioed' across three electrodes. If the ratio of the unknown matches the ratio of the standard, this can help to confirm the identity of the compound. Any difference in the ratio signifies that either there is some contamination of this compound with another unknown compound or that this compound is not the same as the standard compound (Svendsen, 1993).

2.10 Mechanism of Ageing and Ageing Related Theories

Aging is characterized as a progressive decline in biological functions with time. The most prominent characteristics of aging are a progressive decrease in physiological capacity, a reduced ability to respond adaptively to environmental stimuli, an increased susceptibility to diseases, and increased mortality (Farooqui and Farooqui, 2010). Ageing increases the susceptibility to the onset of age related diseases such as Alzheimer's, Parkinson's, diabetes, cancer, autoimmune, and cardiovascular diseases (Kirkwood and Austad, 2000). Aging is associated with several changes in structure and function of different organs and tissues. The liver is the major organ responsible for a number of physiological processes that are crucial to sustaining life. Liver ageing may disturb the liver functions.

In the aging of heart adverse metabolic alterations with complex changes in cardiovascular structure and function take place which can accelerate vascular degenerative diseases (Zhu et al. 2001; Ferrari et al., 2003). In addition, the incidence and prevalence of coronary disease, hypertension, heart failure and stroke increase exponentially with advanced age (Forman et al., 2010). Cardiovascular disease is the leading cause of death and illness in developed countries. It is estimated that by the year 2020 almost 40% of all deaths will be due to cardiovascular disease (Chung et al., 2009).

Many ageing theories were proposed including genome instability theory which postulates that genomic modifications in ribosomal RNA, mitochondrial DNA play an important role in ageing. Telomerase theory is based on the shortening of telomeres due to lack of telomerase activity with ageing. Also it is stated that cell death triggers ageing process (Johnson et al., 1999).

Among them the most accepted theories of aging is the free radical theory, which postulates that aging is the result of the accumulation of cellular and tissue damage caused by the continuous and endogenous generation of reactive oxygen species (ROS) (Sohal, 2002; Harman, 2003; Barja, 2004). Reactive species are produced continuously as a result of normal metabolic processes. At low levels, ROS function as signaling intermediates for the modulation of fundamental cell activities such as growth and adaptation responses.

But if they are produced in excessive amount, ROS contribute to damage in cells and the imbalance between the antioxidant defense system and reactive species is observed. As a result oxidative stress occurs, that contribute to the pathogenesis and progression of age related disorders (Kregel and Zhang, 2007).

Oxidative stress in a physiological setting can be defined as an excessive bioavailability of ROS, which is the net result of an imbalance between production and destruction of ROS (with the latter being influenced by antioxidant defenses) (Hagen, 2003). As a result of a rise in intracellular oxidant concentrations, various cell components are damaged and activation of specific signaling pathways is triggered. Both can influence numerous cellular processes linked to aging and the development of age-related diseases (Finkel and Holbrook, 2000; Valko et al., 2007).

The damage on cell components includes modification of molecules and other cellular components, and is called oxidative damage. Main targets for oxidative damage are proteins, lipids and nucleic acids of living organism as shown in Figure

2.3 and some of their oxidation products are often used as biomarkers of oxidative stress (Blumberg, 2004; Dalle-Donne et al, 2006).




Figure 2.3: Main targets of oxidative stress.

Another theory for ageing is the molecular inflammation hypothesis suggesting that there is link between chronic, low grade inflammation and the pathogenesis of age-related diseases, it is based on dysregulation of the immune system with age and altered redox status during ageing. Both processes lead to activation of a wide variety of inflammatory mediators through overproduced reactive species during aging (Chung et al., 2009).

2.11 Biomarkers of Oxidative Stress

As a biomarker of oxidative stress there are several parameters. But in our study lipid peroxidation, protein oxidation, proteasomal activity, some biomarkers of inflammation and the mRNA expression level of some genes as a member of antioxidant defense system were investigated.

2.11.1 Lipid peroxidation

Lipids are also important targets for oxidation by radical species. Oxidative stress induced peroxidation of membrane lipids leads to alterations in the biological properties of the membrane, such as fluidity, and can lead to inactivation of membrane bound receptors or enzymes, which in turn may impair normal cellular function and increase tissue permeability (Dalle-Donne et al., 2006).

Lipid peroxidation has been extensively studied biomarker of oxidative stress as polyunsaturated fatty acid components of phospholipids in cellular membranes are sensitive to free radical attack. The most prevalent method is the measurement of lipid peroxidation by the thiobarbituric acid reactive substances (TBARS) assay, in which a color is produced by the reaction of aldehydic groups on products e.g., malondialdehyde (MDA). Besides F2-isoprostanes, and 4-hydroxy-2-nonenol (4-HNE), which arose from free radical initiated oxidative damage of polyunsaturated fatty acids can be measured as a biomarker (Bokov et al., 2004).

2.11.2 Nucleic acid oxidation

The hydroxyl radical is known to react with DNA molecule, damaging the purine and pyrimidine bases and also the deoxyribose backbone. The most extensively studied DNA lesion is the formation of 8-OH-G. Modification of genetic material due to oxidative damage is known to be the first step involved in mutagenesis, carcinogenesis, and ageing. The level of 8-oxo-2-deoxyguanosine (oxo8dG) in DNA has been consistently used as a measure of oxidative damage to DNA in ageing (Valko et al., 2007).

2.11.3 Protein oxidation

As proteins are major components of biological systems and play an important role in many cellular functions such as signal transduction, cellular transport systems, age-related increase in oxidative damage to proteins could be physiologically important. Almost all amino acids in proteins are potential targets for oxidation by ROS (Bokov et al., 2004).

Exposure of proteins to ROS may alter every level of protein structure, causing physical changes in protein structure. Oxidative damage to proteins lead to peptide backbone cleavage, cross-linking, and/or modification of the side chain of amino acids (Stadtman and Berlett, 1997; Davies, 1999). These oxidative modifications, inhibit enzymatic and binding activities, increase susceptibility to aggregation and proteolysis. In addition, accumulation of the modified proteins disrupts cellular

function either by loss of catalytic and structural integrity or by interruption of regulatory pathways.

The most widely studied modification to proteins is the formation of carbonyl derivatives on lysine, arginine, proline, histidine, cysteine, and threonine residues. These biomarkers are useful indicators of redox status and have been shown to accumulate during ageing and age-related disease in variety of organisms (Stadtman and Berlett, 1997).

2.11.4 Decreased proteasomal activity

Proteasome is an intracellular proteolytic complex involved in a variety of cellular processes, including the removal of altered forms of protein, growth and differentiation, immune and stress response, signal transduction, aging and prevention of apoptosis (Zeng et al., 2005). The 20S proteasome is the proteolytic core of the proteasomal system. It is composed of two sets of 14 different subunits and has a molecular mass of 670-700 kDa. The subunits form a cylinder of four rings, each containing seven subunits. It has three proteolytic centers showing either peptidyl-glutamyl peptide hydrolyzing (β -1), trypsin-like (β -2), or chymotrypsin-like activity (β -5) based on the proteolysis of peptide substrates for cleavage after acidic basic and hydrophobic residues respectively (Breusing and Grune, 2008). It has been reported that proteasomal activity decreases with age (Viteri et al., 2004; Breusing et al., 2009).

2.11.5 Increased inflammation

During ageing reactive species are overproduced due to oxidative stress-induced redox imbalance. The main cause of age-related redox imbalance is the weakened anti-oxidative defense system which activates the immune systems (Brod, 2000). In human studies, circulating levels of pro-inflammatory cytokines such as TNF- α , IL-6, and IL-1 β , are well-recognized biomarkers of inflammation. Also inflammatory cell (neutrophils, monocytes) counts increase during aging (Bruunsgaard, 2006). Furthermore, aging is associated with increased levels of C-reactive protein (CRP) and serum amyloid P (SAP) proteins (Bruunsgaard et al., 1999). Chronic inflammation is a major underlying condition of many age-related diseases, such as atherosclerosis, cancer, diabetes, osteoporosis, dementia, vascular diseases, obesity and metabolic syndrome (Yu and Chung, 2006).

2.12 Protection from Detrimental Effects of Oxidative Stress

To protect the body from detrimental effects of oxidative stress, the members of antioxidant defense system, including both non-enzymatic (tocopherols, vitamin C, glutathione, etc.) and enzymatic antioxidants (superoxide dismutases SOD, glutathione peroxidases, catalase, etc.) play important roles. In our study vitamin C and vitamin E, glutathione, uric acid, paraoxonase family members (PON-1 and PON-2) and Phase II enzymes were investigated.

2.12.1 Antioxidant defense system

2.12.1.1 Vitamin C (Ascorbic acid)

Ascorbic acid is an essential micronutrient in primates which exhibits some of its biological activity due to its free radical scavenging activity. It is a water soluble chain breaking radical scavenger and recycles plasma membrane α -tocopherol via the reduction of the α -tocopheroxyl radical (May, 2000). Ascorbic acid is transported into mammalian cells either by the Sodium-Vitamin C Transporters, SVCT1 and SVCT2, which cause Na^+ -dependent uptake of L-ascorbate or glucose transporter (GLUT). Unlike humans, primates and guinea pigs, rodents such as laboratory mice exhibit an intact endogenous ascorbic acid synthesis since they express the rate limiting enzyme of vitamin C synthesis which is gulonolactone oxidase, the last enzyme in ascorbic acid synthesis. It has been reported that primary role of SVCT1 is maintenance of the whole-body homeostasis, through dietary absorption and renal reabsorption, while SVCT2 is crucial for ascorbic acid uptake in metabolically active and specialized tissues, such as eye, brain, skeletal muscle and heart (Savini et al., 2008). Beside Kuo et al. (2004) showed that SVCTs appear to function independently of each other, as SVCT1 expression and the ascorbic acid concentration in SVCT1-predominant organs such as liver.

2.12.1.2 Vitamin E (α -tocopherol)

Vitamin E is a generic name for all substances exerting the biological activity of α -Tocopherol. The eight natural existing Vitamin E-compounds (α -, β -, δ -, γ -Tocopherol and α , β , δ -, γ -Tocotrienol) are only generated in plants and therefore they must be supplied to the body. α -Tocopherol the most abundant form of dietary vitamin E, and the major lipid soluble chain-breaking antioxidant in vivo, protecting membranes from oxidative damage. It has been recognized as a potent inhibitor of lipid peroxidation in cellular membrane through the scavenging of peroxy radicals

forming tocoperoxyl radicals (Burton et al., 1982). Apart from its antioxidant activity, it has anti-cancer, anti-atherosclerotic, and neuroprotective effects. Beside antioxidant gene-regulatory activities of α -tocopherol have also been described.

Lipoproteins take part in α -tocopherol transport and its metabolism is closely linked to lipid and lipoprotein metabolism and common proteins are responsible for vitamin E metabolism. Vitamin E is absorbed in the intestine and incorporated in chylomicrons. After entering the circulation remnant chylomicron-associated vitamin E is delivered to the liver, where the α -tocopherol form is selectively incorporated into very low-density lipoproteins (VLDL) by the hepatic α -tocopherol transfer protein (TTP) and resecreted into the blood stream. Subsequently, VLDL and low-density lipoproteins (LDL) exchange α -tocopherol with high-density lipoproteins (HDL) (Kayden et al., 1993; Traber and Arai, 1999). α -tocopherol transfer from plasma to cells occurs by lipid transport between lipoprotein and cells. Uptake is facilitated by a) lipid transfer proteins and lipases (lipoprotein lipase-LPL), b) receptor-mediated lipoprotein endocytosis (LDL Receptor-LDLR; LDL receptor-related protein 1-LRP1), c) selective lipid uptake (scavenger receptor class B type I- SR-BI). Intracellular α -tocopherol is subjected to microsomal degradation by the cytochrome P450 family 3A (CYP3A) enzyme subfamily. Delivery of α -tocopherol to liver is facilitated by the activity of the α -tocopherol transport protein (α -TTP) and ATP binding cassette A1 (ABCA1). Excessive amounts of α -tocopherol may be also excluded via the multidrug resistance transporter 1 (MDR1) (Huebbe et al., 2009).

Another compound in vitamin E metabolism is Vitamin C (ascorbate) which regenerate the membrane bound tocoperoxyl radical to tocopherol. In turn, the ascorbyl radical may be regenerated to ascorbate by thiol or polyphenol antioxidants within the so-called antioxidant network (Packer et al., 2001).

2.12.1.3 Glutathione (GSH)

GSH is a tripeptide containing glutamic acid, cysteine and glycine and the predominant non-protein thiol. It has vital functions in the cell, including; the transport of amino acids, maintenance of proteins in a reduced state and protection of cells against reactive oxygen species, drugs or heavy metal ions, detoxifying electrophiles; maintaining of the cellular redox status, maintaining the essential thiol status of proteins by preventing oxidation of -SH groups or by reducing disulfide bonds induced by oxidant stress; scavenging free radicals; providing a reservoir for cysteine; and modulating critical cellular processes such as DNA synthesis, microtubular-related processes, and immune function (Lu, 1999). GSH exists

primarily in two redox forms, i.e., reduced GSH and glutathione disulfide (GSSG, the oxidized form). The GSSG/GSH ratio is considered to be the cellular redox buffer and it can serve as a good indicator of the cellular redox state of the cell. Redox state is a term that has historically been used to describe the ratio of the interconvertible oxidized and reduced form of a specific redox couple. It describes the redox environment of the cell. Redox couples in cells are, responsive to electron flow that changes in the reducing/oxidizing environment (Schafer and Buettner, 2001).

Also amino thiols such as methionine, cysteine, homocysteine take part in the synthesis of GSH, thus in redox state. The amino thiol metabolism that leads to GSH synthesis is given in Figure 2.4.




Figure 2.4 : Amino thiol metabolism and GSH synthesis 1) Methionine adenosyl transferase 2) transmethylation reactions 3) S- adenosyl homocystein hydrolase 4) cystathionine β-synthase 5) γ-cystathionase 6) γ-glutamyl cysteine synthetase 7) GSH synthetase.

2.12.1.4 Uric acid

Uric acid is the final product of purine metabolism in human. In blood, it is a powerful antioxidant and is a scavenger of singlet oxygen, hydroxyl radical, and peroxynitrite. In humans, approximately one half of the antioxidant capacity of plasma comes from uric acid (Becker et al., 1993). Recently it has been stated that uric acid increases

the cysteine uptake and GSH synthesis in brain (Aoyama et al., 2011) and in liver (Allameh et al., 2008).

On the other hand it has been shown that uric acid is an independent risk factor for cardiovascular disease and metabolic abnormalities (Bonora et al., 1996). Increased uric acid levels have been associated with the presence of the metabolic syndrome (Choi et al., 2007). Indeed, hyperuricemia often precedes the development of hyperinsulinemia, obesity, and diabetes (Johnson et al., 2009; Dehghan et al., 2008). It has been reported that uric acid may also induce inflammatory responses, elevate serum triglyceride and cholesterol concentrations, blood glucose levels (Bonora et al., 1996; Lin et al., 2007).

2.12.1.5 Paraoxonase (PON)

Paraoxonases (PON1, PON2, and PON3) are a unique family of calcium-dependent hydrolases, with enzymatic activities toward a broad range of substrates (lactones, thiolactones, carbonates, esters, and phosphotriesters). The activity of PON1 and PON3 are HDL associated proteins that have been reported to contribute to the anti-atherogenic effects of HDL (Ng et al., 2006). In contrast to PON1 and PON3, PON2 does not associate with HDL in the circulation and it is an intracellular membrane-associated protein. PON1 is widely expressed in liver and PON3 in liver and kidney. Beside PON2 is widely expressed in a number of tissues with the highest expression in the heart, brain, liver, lungs, and testis (Devarajan et al., 2011).

PON1 is known as a longevity gene (Lescai et al., 2009) and loss of PON1 activity with aging has been described in humans (Seres et al., 2004; Jaouad et al., 2006). PON-1 is an enzyme that first synthesized in the liver and subsequently released into the blood (Thomás-Moyá et al., 2006). It protects LDL and HDL particles from oxidative stress (Aviram et al., 1998) and exerts anti-inflammatory properties (Dullaart et al., 2009). Low serum PON1 activity has been found in numerous pathological conditions, such as cardiovascular disease, type1 and 2 diabetes, hypercholesterolemia, metabolic syndrome and renal failure (Mackness et al., 2003; Lescai et al., 2009). It is suggested that PON2 may protect against oxidative stress (Ng et al., 2005; Reddy et al., 2008), atherosclerosis, and foam cell formation (Rosenblat and Aviram, 2009; Ng et al., 2006) it may act as a cellular antioxidant, play an antiatherogenic role (Rosenblat et al., 2006; Ng et al., 2006). It is reported that PON-2 deficient mice show atherosclerotic lesions with enhanced mitochondrial dysfunction (Devarajan et al. 2011), accumulation of triglycerides in macrophages, liver, aortas (Rosenblat et al., 2009; Meilin et al., 2010), increased atherosclerosis

(Ng et al., 2006). Decreased PON2 mRNA and protein levels were demonstrated in human atherosclerotic plaque areas as compared with plaque-adjacent region and normal artery (Fortunato et al., 2008).

2.12.2 Phase II metabolizing enzymes

Phase II metabolizing enzymes are the subgroup of drug metabolizing enzymes, which play central roles in the metabolism, elimination and/or detoxification of xenobiotics or exogenous compounds and drugs introduced into the human body. They protect the body against the potential harmful exposure to xenobiotics from the environment as well as certain endobiotics. In order to minimize the potential injury caused by these compounds, most of the tissues and organs are well equipped with these enzymes (Xu et al., 2005). Some of the Phase II metabolizing enzymes are GST, γ -GCS, NOQ-1, and HO-1.

2.12.2.1 NADPH quinone oxidoreductase (NOQ-1)

NOQ-1 is an enzyme that detoxifies the cycling quinones via reduction thus, it protects cellular membranes against oxidative damage. As a result of reduction, stable hydroquinones are formed that can be readily conjugated and excreted. NOQ-1 is known to have antioxidant properties. It takes part in the regeneration of antioxidant forms of ubiquinone (coenzyme Q) and Vitamin E with the reduction mechanism. It effectively catalyzes the reduction of α -tocopherolquinone (a product of vitamin E) to α -tocopherolhydroquinone, a compound which has antioxidant properties (Ross et al., 2000).

2.12.2.2 Glutathione-S-transferase (GST)

GST is one of the detoxification enzymes that catalyze the conjugation of both endogenous and exogenous compounds with electrophilic moiety, toxic or carcinogenic substrates that cause oxidative stress, with GSH. It has been postulated that GST activity might result in more efficient elimination of carcinogens and ultimately lead to cancer prevention GST catalyzes the addition of the glutathione moiety to a great variety of acceptor molecules. This conjugation is usually considered as a detoxification procedure because conjugate is degraded by the enzymes of the γ -glutamyl cycle (Wilkinson et al., 1997).

2.12.2.3 Gamma glutamyl cysteinyl synthetase (γ -GCS)

γ -GCS is the rate limiting enzyme in GSH synthesis and its activity is a major determinant of intracellular GSH homeostasis. It is composed of two subunits,

namely catalytic and regulatory subunits. It has been suggested that, without the regulatory subunit, the catalytic subunit may not have any activity under physiological.

2.12.2.4 Hemeoxygenase-1 (HO-1)

HO-1 is the rate limiting enzyme in the oxidative degradation of heme to biliverdin, free iron and carbon monoxide. Free heme can be quite cytotoxic, it can rapidly intercalate with membranes and cause severe damage. Exposure of microsomal membrane fractions to free heme in the presence of H_2O_2 leads to a dose-dependent peroxidation of unsaturated lipids in vitro. Heme also catalyzes the protein peroxidation of susceptible groups in membranes (Morita et al., 2005). HO-1 is well known for its antioxidant, anti-inflammatory, antiapoptotic and cytoprotective functions (Ferris et al., 1999; Petrache et al., 2000; Idriss et al., 2008). It may have crucial roles in the pathogenesis of cardiovascular diseases, angiogenesis, diabetes mellitus and cerebrovascular diseases (Idriss et al., 2008).

All of these enzymes, GST, γ -GCS, NQO-1, PON, HO-1 are regulated under the control of nuclear factor erythroid 2-related factor 2 (Nrf2) transcription factor. Nrf2 is an important molecular switch which orchestrates the gene expression of antioxidant and phase 2 enzymes, centrally involved in the ageing process (Lewis et al., 2010; Sykietis et al., 2011). Nrf2 is a basic leucine zipper transcription factor that binds to the antioxidant response element (ARE) in the promoter region of many adaptive genes, such as GST, γ -GCS, NQO1. Under basal conditions, Nrf2 is sequestered in the cytosol. Upon activation, Nrf2 is released, translocates into the nucleus where it heterodimerizes, binds to the ARE and subsequently increases Nrf2 target gene expression (Jaiswal, 2004).

Researchers have demonstrated the role of several genes may modulate aging and influence longevity. The aim is to find the ways for healthy longevity rather than enhancing human longevity (Farooqui and Farooqui, 2010). This may be mediated through modulating age-related genes via phytochemicals, dietary interventions. It has been recently shown that hydroxytyrosol induces antioxidant/detoxifiant enzymes and Nrf2 translocation in hepatocytes (Martin et al., 2010). Furthermore hydroxytyrosol induced cytoprotection against oxidative injury in vascular endothelial cells via Nrf2 dependent signal transduction pathway (Zrelli et al., 2011). Little is known about the role of olive oil phenolics on Nrf2 dependent gene expression in vivo.

Apart from Nrf2 and antioxidant enzymes, in recent years many studies have focused on SIRT1 gene, a class III deacetylase, that can modulate numerous longevity promoting cellular functions, endothelial function, lipid metabolism, cell cycling, gene expression, cardioprotection and aging. It decreases glucose levels, improves insulin sensitivity, decreases production of ROS, modulates DNA repair, and apoptosis (Longo and Kennedy, 2006). Furthermore activation of SIRT1 by polyphenols may be beneficial for regulation of oxidative stress, ageing and metabolism (Chung et al., 2010).

There is a great interest in the development of anti-ageing therapies, protecting from the deleterious effects of ageing, and antioxidants have been widely investigated in this respect. Olive oil, a food naturally containing high amounts of phenolic compounds, may be an alternative treatment. To understand the effect of olive oil phenolics on ageing in vivo studies are needed in human ageing models.

2.13 Models of Human Ageing and the Senescence-Accelerated Mouse Prone (SAMP)

In order to study human ageing yeasts, worms, flies, cells, rodents (aged rats and mice) are used. Among these models, senescence-accelerated mouse (SAM) is an animal model of human aging that is widely used in aging research.

Rodent models may be used to understand the cellular and molecular mechanisms of age-dependent degeneration, and to develop dietary interventions for healthy ageing (Chiba et al., 2009). The senescence-accelerated mouse (SAM) is a model of accelerated senescence that was established through phenotypic selection from a common genetic pool of AKR/J strain of mice. In 1975, certain littermates of AKR/J mice were noticed to become senile at an early age and had a shorter life span. Five of these litters with early senescence were selected as the progenitors of the senescence-accelerated-prone mice (SAMP). Three litters with normal aging process were also selected as the progenitors of senescence-accelerated-resistant mice (SAMR). Thereafter, selective inbreeding was applied based on the degree of senescence, the lifespan, and the age-associated pathologic phenotypes (Butterfield and Poon, 2005).

Among the two strains of SAMs, namely senescence-accelerated prone mice (SAMP) and senescence-accelerated resistant mice (SAMR), SAMR strain serves as the control strain that ages normally. Compared with the SAMR strain, the SAMP strain has higher oxidative stress (Sato et al., 1996, Rebrin et al., 2005; Álvarez-García, et al., 2006), elevated biomarkers of lipid (Matsugo et al., 2000; Yasui et al.,

2003; Rodriguez et al., 2008) and protein oxidation (Nabeshi et al., 2006; Okatani et al., 2002), inflammation (Tha et al., 2000), mitochondrial dysfunction (Rodriguez et al., 2008; Carratero et al., 2009), deteriorated antioxidative defense systems (Boldyrev et al., 2001), extensive liver dysfunction (Ye et al., 2004), an early onset of atherogenesis (Fenton et al., 2004), ultimately leading to a shorter lifespan (Takeda et al., 1997). These properties render SAMP8 mice as a suitable rodent model for experimental ageing research (Satoh et al., 2004).

It has been reported that antioxidant compounds would have beneficial effects as anti-ageing compounds, and several studies have been carried out to ascertain this in SAMP mice. Successful results were obtained by melatonin (Carratero et al. 2009, Garcia et al., 2010, Cuesta et al., 2010), growth hormone (Forman et al., 2010), Oligonol (Tomobe et al., 2007), Pycnogenol (Liu et al., 1998), green tea catechins (Unno et al., 2004; Kichido et al., 2007), mulberry extract (Shih et al., 2009), proanthocyanidins (Gong et al., 2008), lipoic acid and N-acetylcysteine (Farr et al., 2003) as well as caloric restriction (Komatsu et al., 2008; Tajés et al., 2010) and exercise in combination with resveratrol (Murase et al., 2009). Most of these studies focused on brain ageing but there is a need to research cardiological ageing that lead to atherosclerosis and coroner heart diseases, liver ageing and associated pathophysiological processes in SAMP mice.

2.14 In Vivo Studies Related to Effects of Olive Oil on Ageing

Studies related to effects of olive oil on ageing are scarce. In the literature there are a few studies that are performed with aged rats. To our knowledge there is no study investigating the effect of olive oil on aging related disorders in SAMP8 mice but studies are warranted in these mice. Pitozzi et al. (2010) treated the aged rats with high fat feed containing phenol rich extra virgin olive oil and examined the parameters related to oxidative stress. At the end of the study DNA damage, lipid peroxidation, GSH and GSSG levels, and glutathion peroxidase, superoxide dismutase (SOD) and xanthan oxidase (XO) activity in the cerebral cortex, were not different among the groups. Also extra virgin olive oil phenols did not exert protective actions on motor and cognitive functions of the ageing brain, but it is reported that they might have an anxiety-lowering effect associated with decreased glutathione reductase activity and expression in the brain. In the similar study the endogenous levels of DNA breaks, DNA oxidized bases and TBARS in the liver did not differ among the treatment groups of aged rats that fed corn oil, olive oil with high phenolic content and olive oil with low phenolic content. However, a significant

increase in SOD activity and decrease in XO activity were found in the rats fed with high phenolic olive oil. In that group a decrease in DNA oxidative damage in blood cells and plasma TBARS levels, lower incidence of hypophyseal tumors and ulcerative dermatitis were also observed that increased with aging (Jacomelli et al., 2010). In both studies it is reported that the high calorie diet containing large amounts of fat overwhelmed the possible beneficial effect of olive oil phenols. Quiles et al. (2004) have investigated the effects of lifelong olive oil feeding in rats, and found increased antioxidant capacity, reduced DNA damage and improved plasma lipid profile as compared to sunflower oil. Bello et al. (2006) reported a regulatory effect of olive oil on hepatic apoptotic pathways in aged rats with increased caspase-9 and caspase-3 activities as compared to rats fed with sunflower oil. Olive oil also decreased Bcl-2 levels, increased membrane bound Bax and decreased plasma membrane Mg^{2+} -dependent neutral sphingomyelinase. The higher CoQ levels found in membranes from the olive oil group in aged rats was also contributed to a better antioxidant protection.

2.15 Aim of the Study

Olive oil phenolics are well known for its antioxidant activities. But their role in metabolic pathways, their effect on transcription factors, gene expressions need to be investigated. In this study it is aimed to study the effect of diet rich in olive oil phenolic compounds on age-related changes in heart and liver tissues of SAMP mice. As oxidative stress and inflammation play an important role in the pathogenesis of ageing and age-related disorders, the parameters regarding these factors were investigated. In both tissues Nrf2 and the genes that are controlled under that transcription factor was the main focus. Their amounts were determined both in mRNA level and protein level. Furthermore the level of antioxidant vitamins (vitamin C and vitamin E) was measured. As a biomarker of oxidative stress, lipid oxidation was determined by TBARS assay, protein oxidation was determined by measuring protein carbonyl levels. Also proteasomal activity of different subunits in the proteasomal system was measured.

Another aim of that study is to develop and validate a method with the sensitive electrochemical detector for the quantification of olive oil phenolics in oils

3. MATERIALS AND METHOD

3.1 Chemicals and Reagents

HPLC grade methanol was obtained from J.T. Baker (Deventer, The Netherlands) and acetonitrile was obtained from Sigma Aldrich (Steinheim, Germany). Standards of tyrosol (CAS no. 501-94-0), vanillic acid (CAS no. 121-34-6), caffeic acid (CAS no. 331-39-5), p-coumaric acid (CAS no. 501-98-4), and ferulic acid (CAS no. 1135-24-6) were purchased from Sigma Aldrich (Schnelldorf, Germany). Oleuropein (CAS no. 32619-42-4) and hydroxytyrosol (CAS no. 10597-60-1) were supplied by Extrasynthese (Genay Cedex, France) and pinoresinol by Separation Research (Turku, Finland). Reference standards of all analyzed compounds were HPLC grade with purity higher than 98% (except oleuropein, >90%). Stock solutions were prepared in methanol and stored at -20°C. HPLC grade 60% perchloric acid was obtained from Fisher Scientific (Leicestershire, UK).

For the protein carbonyl quantification potassium chloride, mono potassium phosphate, sodium chloride, sodium phosphate dibasic dihydrate, disodium phosphate, monosodium phosphate and guanidine hydrochloride, citric acid, sulfuric acid, hydrogen chloride, sodium hydroxide, bovine serum albumin and Roti Quant® were purchased from Carl Roth GmbH (Karlsruhe, Germany). Tween 20, 2,4 dinitrophenylhydrazine, biotin-conjugated rabbit IgG polyclonal antibody raised against a DNP conjugate of keyhole limpet hemocyanin (anti-DNP), streptavidinbiotinylated horseradish peroxidase, and o-phenylenediamine were from Sigma-Aldrich Chemie GmbH (Steinheim, Germany). Hydrogen peroxide was obtained from Merck KGaA (Darmstadt, Germany).

3.2 Sampling of Olive Oil

In total 57 olive oils from Italy, Spain, France, Turkey, Portugal, Australian, New Zealand, South Africa which were directly collected from producers or supermarkets were analysed in terms of their phenolic content. The oil with the lowest (Coopérative Oléicole du Canton d'Ollioules, France) and highest total phenolic content (Fratelli Ferrara s.a.s, Italy) was then chosen for the feeding study in mice.

3.3 Preparation of Olive Oil Extracts

The phenolic fraction of the oil was obtained by solid-phase extraction (SPE) using the diol-bonded SPE cartridges (Varian, Bond Elut Diol 500 mg/3 mL) according to the procedure of Mateos et al. (2001). As a modification oil sample of 1.5 g was mixed with 1.5 mL hexane. The phenolic extract was redissolved in 1 mL methanol. Samples were filtered through 0.2- μ m PTFE Membrane filter (Phenomenex, USA) into the vial.

3.4 Spectrophotometric Determination of Total Phenolic Content

The total phenolic content of the extracts was determined according to the Folin-Ciocalteu spectrophotometric method (Singleton et al., 1999). Diluted Folin-Ciocalteu reagent and 20% sodium carbonate solution were added to the phenolic extract. The solution was left in the dark for 45 min. The absorbance of each sample as well as gallic acid standards was determined at 750 nm. The results were expressed as mg gallic acid/kg of oil.

3.5 HPLC Analysis of Olive Oil Phenolics

The HPLC analyses were carried out on a Jasco system (Jasco GmbH Deutschland, Gross-Umstadt, Germany) consisting of a pump (PU-2085), autosampler (XLC-3059AS) with integrated column heater, while detection was carried out on 8-channel ESA Model 5600A CoulArray Detector (ESA Inc., Chelmsford, MA, USA). For analyte detection, increasing potentials of +250, +400, +500 and +750 mV were applied on channels. Separation of compounds was carried out on a fused core type column (Kinetex C18, 2.6 μ m, 100 $\times$ 4.6mm, Phenomenex, USA). Column temperature was maintained at 40°C and flow rate was 1.25 mL/min. Mobile phase was composed of water (solvent A), acetonitrile (solvent B) and methanol (solvent C) each containing 60mM LiClO₄. The pH was adjusted to 3.1 with perchloric acid, and the eluent was vacuum-filtered through a 0.2 μ m hydrophilic polypropylene membrane filter (Pall Corporation, Michigan, USA). Separation was achieved by elution gradient using an initial composition of 96% A 3% B 1% C. The composition changed to 85% A 10% B 5% C at 4 min, 82.5% A 11.5% B 6% C at 6 min, 65% A 24% B 11% C at 15 min and finally increased to 85% B 15% C at 21 min and maintained for 5 min. Initial conditions were reached in 5 min. The flow rate was 1.25 mL/min and injection volume was 10 μ L.

3.6 Method Validation

The analytical method for the determination of olive oil phenolics was validated according to the requirements of FDA (FDA, 2001) in terms of linearity, selectivity, lower limit of quantification and detection, accuracy, precision, recovery, stock solution stability, short term stability, freeze thaw stability and post preparative stability.

3.6.1 Selectivity

For evaluation of selectivity, phenolic compound free olive oil (Sigma Aldrich, Steinheim, Germany) extracted and analyzed to check for possible interference with the retention time of the analytes.

3.6.2 Linearity, lower limit of detection (LLOD) and lower limit of quantification (LLOQ)

The linearity range of the detector response was performed by serial dilution of a stock solution of the studied compounds. Stock solutions were prepared in methanol and further diluted by H₂O/methanol (50:50). Regression parameters were calculated for calibration curves. All the calibrations curves were linear over the range of study and were calculated by using seven points at different concentrations.

The lower limit of quantification is defined as the concentration of the lowest standard in the analytical run which is quantified with a deviation of the actual concentration and a coefficient of variation of precision of less than 20%. The limit of detection (LOD) was the lowest detectable analyte concentration at the retention time where no peak of the analyte was observed.

3.6.3 Accuracy

The accuracy of the method was determined by five replicate analyses of quality control samples, which were obtained by spiking the phenolic free olive oil with known amounts of olive oil phenolic standards at three concentrations (low, medium, high). The olive oil was extracted and the calculated analyte concentrations of spiked oil samples were compared with the theoretical concentration values. The highest amount of phenolics added to the olive oil were determined according to the literature in order to study with realistic values. This was diluted by 4 for the middle concentration and by 20 for the low concentration.

3.6.4 Precision

The precision of the method was determined by five replicate analyses of quality control samples, which were obtained by spiking the phenolic free olive oil with known amounts of olive oil phenolic standards at three concentrations (low, medium, high) on three consecutive days. The olive oil was extracted and the calculated analyte concentrations of spiked oil samples were compared with the theoretical concentration values within the same day (inter-day) and from day-to-day (intra-day) for three consecutive days. Values should not deviate by more than 15 % (high and medium concentrations) or 20 % (low concentration). The highest amount of phenolics added to the olive oil were determined according to the literature in order to study with realistic values. This was diluted by 4 for the middle concentration and by 20 for the low concentration.

3.6.5 Recovery

The recovery of phenolics calculated by analyzing five replicates for each concentration (low, middle, high) and comparing the detector responses with those of calibration standards with identical concentrations.

3.6.6 Stock solution stability

Inject stock solution (if necessary prepare appropriate dilution) into HPLC and store aliquots of the stock at -20°C and room temperature for 24 h. Compare initial detector response to that after 24 h storage. The stability of stock solution were determined for three concentration levels (low, medium and high) after storage at room temperature, 4°C, -20°C, -80°C for 24 h.

3.6.7 Short term stability

Five aliquots of spiked oil samples were prepared at 3 concentrations (low, medium and high). The first aliquot was analysed at 0h and remaining samples were stored at room temperature, 4°C, -20°C, -80°C for 24 h. To determine stability initial values were compared to those obtained after storage at different temperatures.

3.6.8 Freeze thaw stability

Three aliquots of spiked oil samples were prepared at 2 concentrations (low and high). Oil samples were analysed at 0h and the remaining three aliquots were frozen at -80°C for 24 h, thawed at room temperature and frozen again for 12-24 h at -80°C. The freeze-thaw-cycle were repeated two additional times. The stability was

determined by comparing initial values to those obtained after repeated freezing-thawing.

3.6.9 Post preparative stability

Three samples of spiked olive oil with low and high concentrations were prepared in order to determine the stability of processed samples in the autosampler. Samples were injected from the same HPLC vial at 0h and compared with the injections at 6h intervals (6h, 12h and 18h). The cups of HPLC vial caps were replaced after each injection to minimize solvent loss.

3.7 FRAP and TEAC Assays

The FRAP assay was conducted according to Benzie and Strain (1996), using ascorbic acid as a reference. The TEAC assay was conducted as described by Miller et al. (1996) using Trolox as a reference.

3.8 Animals and Study Design

The animal experiment was performed according to German animal welfare laws and regulations and with permission of the appropriate authorities. Twenty two female SAMP8, aged 9-10 weeks were obtained from Harlan Winkelmann GmbH (Borchen, Germany). The mice were housed in groups in type II polypropylene cages equipped with softwood bedding, a water bottle, a mouse house and a table tennis ball in a climate-controlled room (temperature, $22 \pm 2^{\circ}\text{C}$; humidity, $55 \pm 5\%$) with a 12 h light/dark cycle. One of the SAMP8 mouse died during the experiment. Mice were fed with pelletized Western type diet (Altromin Spezialfutter GmbH & Co. KG, Lage, Germany) with 0.15 % cholesterol and 20 % fat, in which 10% of fat was from olive oil. Mice were randomly divided into two groups. All animals had free access to feed and water. Food intake was controlled daily and body weights were controlled weekly during the 4.5 month feeding trial. At the end of the experiment the mice were starved before anaesthesia by CO_2 and decapitated. Blood samples were collected in tubes and serum was separated by centrifugation (Eppendorf 5804 R, Rotor F34-6-38, Wesseling-Berzdorf, Germany). Tissues were excised in liquid nitrogen and stored at -80°C until analysed.

3.9 PON1 Arylesterase Activity in Serum

Arylesterase activity was measured by using phenylacetate as an artificial substrate for PON1. Initial rates of hydrolysis were determined spectrophotometrically at 270 nm. The assay mixture included 4 mmol/L of phenylacetate and 1 mmol/L of CaCl₂ in 20 mmol/L of Tris HCl, pH 8.0. Non-enzymatic hydrolysis of phenylacetate was subtracted from the total rate of hydrolysis. One unit of arylesterase activity is equal to 1 µmol of phenylacetate hydrolyzed per minute per milliliter (Fuhrman et al., 2006).

3.10 Liver and Heart Tissue Preparation

A ratio of 1:10 homogenates of liver and 1:12 of heart were prepared with ice cold phosphate buffered saline (PBS) buffer (pH 7.4) at 26.000 rpm with a Micra D-8 homogenizer (ART Prozess- & Labortechnik GmbH & Co. KG, Mullheim, Germany). Homogenates were centrifuged at 4,800 *g* at 4°C for 10 min (Eppendorf 5804 R, Rotor F34-6-38, Wesseling-Berzdorf, Germany) and the supernatant was stored at -80°C until further use.

3.11 Lipid Peroxidation

Lipid peroxidation was assayed fluorometrically with Tecan Infinite 200 microplate (Tecan Group Ltd., Crailsheim, Germany) as thiobarbituric acid reactive substances (TBARS) in heart homogenates after protein precipitation with TCA and extraction in 1-butanol. Excitation and emission wavelengths were 520 nm and 560 nm, respectively. Calibration curve was prepared with TEP (1,1,3,3-tetraethoxypropane) as an external standard (Morel et al., 1983).

3.12 Quantification of Protein Carbonyls

Protein carbonyl content was determined in the homogenized heart tissue supernatant according to the ELISA method by Buss et al. (1997) with required modifications. The detection system used an anti-dinitrophenyl rabbit IgG-antiserum (Sigma Aldrich, Steinheim, Germany) as the primary antibody and a monoclonal antirabbit IgG antibody peroxidase conjugate (Sigma Aldrich, Steinheim, Germany) as the secondary antibody. Colour change was induced with *o*-phenylenediamine and H₂O₂.

3.13 Proteasomal Activity

Twenty to forty mg tissue was homogenized in lysis buffer (250 mM sucrose, 25 mM HEPES, 10 mM magnesium chloride, 1 mM EDTA and 1.7 mM DTT) using a homogenizer (ultra-Turrax®) and afterwards centrifuged at 14,000 x g for 30 min. The supernatant was used for determination of protein content using the Bradford assay and for measurement of the proteasomal activity. For proteasomal activity, samples were incubated in 225 mM Tris buffer (pH 7.8), 45 mM potassium chloride, 7.5 mM magnesium acetate, 7.5 mM magnesium chloride and 1 mM DTT. For the peptidyl-glutamyl like-(β 1), trypsin like-(β 2) and chymotrypsin like-(β 5) activity the substrates Z-Leu-Leu-Glu-MCA (Biochem, Boston, USA), Ac-Arg-Leu-Arg-MCA (Biochem, Boston, USA) and N-Succinyl-Leu-Leu-Val-Tyr-MCA (Sigma Aldrich, Steinheim, Germany) were used, respectively. MCA liberation of the substrates was measured with a fluorescence reader at 360 nm excitation and 460 nm emission. Free MCA was utilized as a standard (Breusing and Grune, 2006).

3.14 α -Tocopherol Analysis

Tissue samples (50 mg) were placed in a test tube with screw-cap and 2 mL ethanol (EtOH) containing 1% ascorbic acid (w/v) and 700 μ L H₂O. Three hundred microlitres of saturated potassium hydroxide (KOH) solution was added, closed, vortexed for 10 s and incubated for 30 min at 70°C in a shaking water bath. Thereafter samples were immediately chilled on ice and 50 μ L butylated hydroxytoluene (BHT)-solution (0.1% BHT in EtOH, w/v) and 2 mL n-hexane were added. Test tubes were mixed per hand and centrifuged 5 min at 1500 rpm. 500 μ L of the supernatant was transferred to a clean test tube and dried at room temperature (Savant SpeedVac; Thermo, Langenselbold, Germany). Samples were reconstituted with 150 μ L MeOH: H₂O (98:2, v/v), and stored at 20°C until analysis. For quantification of the tocopherols, 40 μ L of the sample were injected into an HPLC system (Jasco, Gross-Umstadt, Germany; pump PU2080Plus, autosampler AS2057Plus, detector FP2020Plus) and separated on a Waters Spherisorb ODS-2 column (3 μ m, 100x4.6 mm) by isocratic elution at a flow rate of 1.2 mL/min with MeOH: H₂O (98:2, v/v) as mobile phase. The fluorescence detector was set to an excitation wavelength of 290 nm and an emission wavelength of 325 nm. Peaks were recorded and integrated using the chromatography software Jasco ChromPass. The concentrations of α - and γ -tocopherol were quantified using an external standard curve with tocopherols (Tocopherol set, Calbiochem®).

Schwalbach, Germany). The same protocol was applied for the quantification of α -tocopherol in olive oils.

3.15 Quantification of Ascorbic Acid, GSH, GSSG, Uric Acid, Cysteine Homocysteine and Methionine by HPLC with Coulometric Electrochemical Detection

Heart and liver tissue homogenates were prepared as described by Rebrin et al. (2005). The chromatographic separation was carried out with an autosampler 851-AS (Jasco GmbH Deutschland, Gross-Umstadt, Germany) and a pump 510 (Waters, Milford, MA, USA) on a Trentec Reprosil-Pur 120 C18 AQ column (150x4.6 mm, 3 μ m) from Trentec-Analysentechnik (Rutesheim, Germany) at a flow rate of 1 mL/min. The mobile phase for isocratic elution consisted of 50 mM sodium phosphate monobasic and 0.25 % (v/v) acetonitrile, the pH was adjusted to 2.5 with 85 % o-phosphoric acid, and the eluent was vacuum-filtered through a 0.2 μ m hydrophilic polypropylene membrane filter (Pall Corporation, Michigan, USA). The column temperature was maintained at 40°C and the analytes were quantified with an 8-channel ESA Model 5600A CoulArray Detector (ESA Inc., Chelmsford, MA, USA). For analyte detection, increasing potentials of +50, +175, +350, +600, +750 and +825 mV were applied on channels. Serial dilutions were prepared from stock solutions in PBS/10 % m-phosphoric acid. The injection volume of samples and calibration standards was 10 μ L. Each sample was analysed in duplicate.

3.16 RNA Isolation and Real-Time Quantitative RT-PCR

RNA was isolated from liver samples (20-30 mg) using TRIsure lysis reagent (Bioline, Luckenwalde, Germany). Real-time quantitative PCR was performed as a one-step procedure (SensiMix One-step Kit; Quantace, Berlin, Germany) with SybrGreen detection, using the Rotorgene cyclor (Corbett Life Science, Sydney, Australia). Relative mRNA concentrations of genes were quantified by the use of a standard curve. Target gene mRNA concentration was normalized to the mRNA concentration of the housekeeping gene GAPDH. Primers were designed by standard tools (Spidey, Primer3 and NCBI Blast) and purchased from MWG (Ebersberg, Germany). Primer sequences for analyzed genes are summarized in Table 3.1.

Table 3.1: Primers used in real time PCR experiments.

Primers	Sequence	Annealing temperature (°C)
ABCA1	F: CTGCCACCTCCTCAGAGAAA R: GCCATACCGAAACTCGTTCA	55
CRP	F: AGATCCCAGCAGCATCCATA R: CAGTGGCTTCTTTGACTCTGC	59
CYP3A	F: CAAGGAGATGTTCCCTGTCA R: CTGGTGATCACATCCATGCT	55
GAPDH	F: GACAGGATGCAGAAGAGATTACT R: TGATCCACATCTGCTGGAAGGT	55
γ-GCS	F: AACACAGACCCAACCCAGAG R: GTCTGGCTGAGAAGCCTTTG	59
Gulo	F: CCAAGATCACACCAACAAGG R: GTCTTGGACATGCTGCTTGA	57
GST	F: TACTTTGATGGCAGGGGAAG R: TCATCCCGTCGATCTCTACC	58
HO-1	F: GAGCCTGAATCGAGCAGAAC R: AGCCTTCTCTGGACACCTGA	59
LCAT	F: TAGCAGGCTACCTGCACACA R: GCCAGCCAGCTTCTGTAGT	59
LDLrec	F: AAGCTGTGACCGATGAGGAG R: CATTGTTCAAGGCCCTGCTTA	55
LPL	F: ATCAACTGGATGGAGGAGGA R: CAGCTGGATCCAAACCAGTA	58
LRP-1	F: AAGCTGTGACCGATGAGGAG R: CATTGTTCAAGGCCCTGCTTA	55
MDR-1	F: GCCAGTATTCTGCCAAGCAT R: CTATGAGCACACCAGCACCA	55
NQO-1	F: TTCTTCTGGCCGATTGAGAGT R: TCCAGACGTTTCTTCCATCC	55
Nrf2	F: GGGGACAGAATCACCATTG R: GATGCAGGCTGACATTCTGA	57
PON1	F: CAGCCTGTCCATCTGTCTCA R: CACCCGTCTCGATTCTTTA	59
PON2	F: ATGGTGGCTCTGAGTTTGCT R: TCCTCAGCTCCAGTTTCGAT	57

Table 3.1 (continued): Primers used in real time PCR experiments.

Primers	Sequence	Annealing temperature (°C)
SAP	F: AAGCTGCTGCTTTGGATGTT R: CATTGTCTCTGCCCTTGACA	55
SIRT1	F: GTCTCCTGTGGGATTCTGA R: ACACAGAGACGGCTGGAACT	57
SRB1	F: GCAAATTTGGCCTGTTTGTT R: GATCTTGCTGAGTCCGTTCC	55
SVCT-1	F: GGGAAAGCCCTCTTCTTTTC R: TGAAGCACGTCAGGTAATGC	57
SVCT-2	F: TATTCCTGGGATTGCAGCAC R: AAAGCACTGGCCTGAAACAG	57
α -TTP	F: GCCAAGAAGATTGCTGCTGT R: GCAGATGAATCCGGTCCTTA	58

3.17 Western Blot Analysis

Liver tissue (20 mg) was homogenized in radio immuno precipitation assay buffer (50mM-Tris-HCl, 150mM-NaCl, 0.5% deoxycholate, 0.1% SDS and 1% NP-40; pH 7.4 with protease-inhibitor cocktail, 1:100; Sigma, St Louis, MO, USA). Lysates were purified by centrifugation (12,000 rpm, 4°C, 20 min) after incubation on ice for 30 min. Total protein concentrations in each lysate were quantified using a BCA Protein Assay kit (Pierce, Rockford, USA). Total protein of the lysates (40 µg per lane) was mixed with loading buffer, denatured at 95°C for 5 min and separated by 12% SDS gel electrophoresis followed by transferring the proteins to a PVDF membrane, which was then blocked for 2h in blocking buffer (3% non-fat milk in Tris-buffered saline, pH 7.4, with 0.05% Tween-20 (TBS/T)). Primary antibodies were diluted in 3% non fat milk (GAPDH (Santa Cruz), 1:500; GST (Abcam), 1:3000; 1:200, NOQ-1 (Santa Cruz), 1:500, HO-1 (Stressgen), 1:1000; Tubulin (Abcam), 1:10000) and the blots were incubated overnight at 4°C. The blots were washed and incubated with the respective IgG secondary antibodies (Santa Cruz Biotechnology, Heidelberg, Germany) in blocking buffer with gentle agitation for 1 h at room temperature. The blots were washed and exposed to Immun-Star Western Chemiluminescent kit (Bio-Rad Laboratories, Hercules, CA, USA) and scanned with a ChemiDoc XRS system (BioRad, Munich, Germany). Digital images were captured and quantified by densitometry using the Quantity-One system (Bio-Rad). Relative concentration of

the proteins was quantified as the ratio between the amount of target proteins and the amount of the housekeeping protein α -tubulin.

3.18 Statistical Analysis

Results are expressed as mean values with SEM. Data were tested for normal distribution and analyzed by *t*-test. Statistical analysis was performed using PASW Statistics 18 (IBM, Chicago, IL, USA). Data were analysed for normality of distribution (Kolmogorow-Smirnov and Shapiro-Wik tests) and equality of variance (Levene's test) before the *t* test for independent samples and in the case of non-parametric data Mann-Whitney U test. Differences were considered significant when the *p* value was ≤ 0.05 .

4. RESULTS AND DISSCUSSION

4.1 Method Validation

For the analysis of selected phenolic compounds of olive oil, the developed method was validated in terms of selectivity, lower limit of quantification and detection, accuracy, precision, recovery, short term stability, freeze thaw stability and post preparative stability. Although all the phenolics can be detected in 16 minutes, the total time of analysis for the proposed method is 30 minutes and equilibration time is 5 minutes. The duration could not be shortened due to the fluctuations caused by the fast gradients, which affect the sensitivity of the detector. As compared to the other methods in the literature, which are longer than 40 minutes, this method gives the opportunity to analyze phenolic compounds in shorter time.

4.1.1 Selectivity

The analytical method was found to be selective for all the analytes and no endogenous substances were found to interfere with the compound separation. Representative chromatogram of the blank olive oil extract is shown in Figure 3 (appendix A).

4.1.2 Linearity, lower limit of detection (LLOD) and lower limit of quantification (LLOQ)

The linearity of the standard curves of the phenolic compounds was detected for ten points measurement. The regression coefficients of all phenolic compounds were over 0.99. The LLOQ and LLOD levels of the phenolics were in ng level. The linear range, LLOQ, LLOD of selected phenolic compounds were given in Table 4.1.

Table 4.1: Linear range, LLOQ, LLOD of selected phenolic compounds for the method validation.

Compound	Linear range ($\mu\text{g/mL}$ - ng/mL)	Regression coefficient	LLOD (ng/mL)	LLOQ (ng/mL)
Tyrosol	16.67-0.3	0.999	0.065	0.3
Hydroxytyrosol	62.5-15.3	0.999	0.062	15.3
Pinoresinol	33.33-2	0.999	0.5	2.0
Oleuropein	222.22-3.4	0.999	0.11	3.4
Caffeic acid	22.22-5.4	0.999	0.3	5.4
Ferulic acid	27.78-3.4	0.999	1.7	3.4
Vanillic acid	40-0.6	0.999	0.15	0.6
p-coumaric acid	22.22-1.4	0.999	0.025	1.4

4.1.3 Accuracy, precision and recovery

The proposed method has good accuracies and reproducibility for all the analytes at all three level of concentrations. The within-day and day-to-day precision have coefficient of variation of <15%. The extraction method was able to produce high recoveries (>72%) for all the analytes as well as the internal standards except middle concentrations of ferulic- and caffeic acid which were 65% and 58% respectively. Further details of the accuracy, precision and extraction recovery are shown in Table 4.2. The highest amount of phenolics added to the olive oil were determined according to the literature in order to study with realistic values. But the recoveries of phenolic acids (caffeic acid, ferulic acid, p-coumaric acid) were very bad (130-160%). Therefore the concentration of these phenolics were increased 10 times.

Table 4.2: Intraday and inter-day recoveries, accuracies and precisions of selected phenolic compounds of olive oil.

	Nominal concentration (mg/kg)	Calculated concentration (mg/kg)	Recovery [%]	Accuracy [%]	Precision [% CV]
Tyrosol					
Intraday (n=5)	1.75	1.64	93.7	6.3	2.56
	8.75	6.50	74.3	25.7	10.15
	35	29.89	85.4	14.6	6.93
Interday	1.75	1.68	95.8	4.2	3.77
	8.75	7.24	82.7	17.3	5.73
	35	30.28	86.5	13.5	6.01
Hydroxytyrosol					
Intraday (n=5)	2.5	1.92	76.6	23.4	3.64
	12.5	8.05	64.4	35.6	10.34
	50	38.65	77.3	22.7	6.69
Interday	2.5	1.99	79.8	20.2	3.78
	12.5	9.14	73.1	26.9	5.40
	50	39.90	79.8	20.2	6.21
Oleuropein					
Intraday (n=5)	0.6	0.61	100.9	-0.9	3.27
	3	2.17	72.2	27.8	13.71
	12	10.45	87.1	12.9	3.24
Interday	0.6	0.56	94.1	5.9	5.75
	3	2.19	72.9	27.1	9.45
	12	9.79	81.6	18.4	10.88
Pinoresinol					
Intraday (n=5)	4.75	4.12	86.8	13.2	3.57
	23.75	17.71	74.6	25.4	12.12
	95	84.55	89.0	11.0	7.43
Interday	4.75	4.30	90.6	9.4	4.24
	23.75	20.38	85.8	14.2	6.72
	95	89.58	94.3	5.7	6.51
Caffeic acid					
Intraday (n=5)	0.75	0.56	74.5	25.5	9
	3.75	2.18	58.1	41.9	11.7
	15	12.58	83.9	16.1	5.44
Interday	0.75	0.57	76.4	23.6	7.68
	3.75	2.76	73.5	26.5	6.16
	15	13.23	88.2	11.8	5.79
Ferulic acid					
Intraday (n=5)	0.75	0.67	89.9	10.1	3.85
	3.75	2.47	65.9	34.1	14.10
	15	13.94	92.9	7.1	4.74
Interday	0.75	0.67	89.8	10.2	4.46
	3.75	2.97	79.3	20.7	6.92
	15	13.75	91.7	8.3	5.55
Vanillic acid					
Intraday (n=5)	0.27	0.26	96.0	4.0	6.24
	1.33	0.97	73.3	26.7	12.43
	5.3	4.46	84.2	15.8	4.53
Interday	0.27	0.26	95.5	4.5	5.49
	1.33	1.05	78.9	21.1	6.55
	5.3	4.46	84.1	15.9	5.78
p-coumaric acid					
Intraday (n=5)	0.75	0.72	96.3	3.7	3.63
	3.75	2.98	79.6	20.4	11.72
	15	14.88	99.2	0.8	4.56
Interday	0.75	0.69	91.6	8.4	4.44
	3.75	3.19	85.2	14.8	6.10
	15	10.80	72.0	5.0	5.37

4.1.4 Stability studies

Stability studies carried out to determine whether any sample deterioration occurred during storage and handling. In terms of stability of stock solutions, the values of high and middle concentrations of phenolic compounds at three different temperatures were slightly different from the freshly prepared standards. However great differences observed for degradation of the low concentrations of hydroxytyrosol at -20°C (-18.31), caffeic acid at room temperature (-23.72) and -20°C (-21.02), ferulic acid at 4°C (-12.36) (Table 4.3).

The analytes were found to be stable after three freeze–thaw cycles as well as in autosampler at room temperature (Table 4.4). Regarding the short term stability and freeze thaw stability, the recovered amount of compounds were >85% for all concentrations at tested four different temperatures. In autosampler stability, greater degradation was observed for the low concentration of caffeic acid and oleuropein with 18.18% and 11.09%, respectively (Table 4.5).

Table 4.3: Stock solution stability of standards after 24 h storage at room temperature (RT), 4°C, and -20°C.

	Stock solution stability [Degradation %]		
	RT	4°C	-20°C
Tyrosol			
Low	-6.77	3.52	5.48
Middle	-2.48	1.64	-0.45
High	-1.68	-0.89	-0.74
Hydroxytyrosol			
Low	-0.85	-6.72	-18.31
Middle	0.62	2.00	0.32
High	-1.31	-0.20	1.52
Oleuropein			
Low	-0.57	-4.96	4.04
Middle	0.71	2.43	0.11
High	-1.21	-0.52	-0.40
Pinoresinol			
Low	-0.64	-4.02	2.37
Middle	2.99	1.83	0.01
High	-1.00	-0.63	-1.20
Caffeic acid			
Low	-23.72	-8.06	-21.02
Middle	2.59	8.84	3.77
High	1.78	1.53	2.0
Ferulic acid			
Low	-8.10	-12.36	3.60
Middle	3.28	2.15	-0.14
High	-1.92	-1.27	-0.94
Vanillic acid			
Low	1.40	-0.57	-1.60
Middle	-0.14	1.96	-0.14
High	-0.56	-0.04	-1.19
p-coumaric acid			
Low	9.90	0.32	2.02
Middle	1.45	2.07	0.26
High	-1.72	-1.19	0.21

Table 4.4: Freeze thaw stability and short term stability of selected phenolic compounds in olive oil at room temperature (RT), 4°C, -20°C, and -80°C.

Nominal Concentration [mg/kg]	Short term stability [Degradation %]				Freeze thaw stability [Degradation %]
	-80°C	-20°C	+4°C	RT	
Tyrosol					
1.75	-3.21	-3.70	-4.05	6.12	10.08
35	-11.49	-10.15	-9.10	-7.46	3.95
Hydroxytyrosol					
2.5	-2.34	-3.07	-1.02	4.39	10.33
50	-12.46	-11.01	-9.0	-5.51	4.21
Oleuropein					
0.6	-3.29	-5.54	-9.65	9.03	-1.56
12	-11.71	-11.07	-9.15	-6.21	7.91
Pinoresinol					
4.75	-6.41	-6.81	-2.71	6.61	9.03
95	-14.45	-13.87	-12.11	-9.57	1.54
Caffeic acid					
0.75	-5.79	-5.47	-5.14	6.82	8.05
15	-13.73	-12.05	-10.11	-6.15	2.25
Ferulic acid					
0.75	-4.75	-5.17	-5.97	7.61	9.28
15	-13.23	-11.83	-7.89	-7.19	2.45
Vanillic acid					
0.27	-5.06	-4.10	-5.63	7.44	11.33
5.3	-14.99	-13.95	-12.40	-9.56	0.10
p-coumaric acid					
0.75	-4.48	-4.89	-4.89	7.54	8.90
15	-11.64	-10.34	-9.10	-6.26	-0.21

Table 4.5: Post preparative stability of selected phenolic compounds in olive oil.

Nominal Concentration [mg/kg]	Post preparative stability [Degradation %]		
	6h	12h	18h
Tyrosol			
1.75	0.00	0.24	0.63
35	0.28	0.70	0.70
Hydroxytyrosol			
2.5	-2.03	-3.93	-0.14
50	-0.42	-0.25	1.38
Oleuropein			
0.6	-5.64	-11.09	1.32
12	0.05	0.21	1.43
Pinoresinol			
4.75	-1.61	-1.85	-0.87
95	0.19	0.00	0.95
Caffeic acid			
0.75	-5.87	-18.18	-5.28
15	-0.51	-1.20	-1.20
Ferulic acid			
0.75	-1.12	-2.77	-0.12
15	0.57	0.57	1.26
Vanillic acid			
0.27	-0.80	-6.97	-0.45
5.3	0.00	-0.26	-0.26
p-coumaric acid			
0.75	-1.04	-2.25	0.69
15	0.59	0.54	1.78

4.2 Application of the Method for the Determination of Selected Phenolic Compounds in Olive Oils

In order to show the applicability of the developed method, the olive oil samples that were collected from different countries were analyzed in terms of selected phenolic compounds. Figure 2 in appendix shows the representative HPLC chromatogram of an olive oil showing the selected eight compounds.

Table 4.6 presents the quantitative results obtained for the analysis of olive oil by HPLC coupled with electrochemical detector. Analyte concentrations were quantified by calibration curves for the phenolic compounds.

Table 4.6: FRAP, TEAC, total phenolic, α -tocopherol, γ -tocopherol and phenolic contents of collected olive oil samples. Values are expressed as mean $\pm$ SD.

Sample	TP	TEAC	FRAP	α -Toc	γ -Toc	TY	HTY	OP	Pino	CA	VA	FA	p-COU
1	532 $\pm$ 10.6	1.02 $\pm$ 0.14	1.28 $\pm$ 0.16	216 $\pm$ 14	9.9 $\pm$ 1.0	20.78 $\pm$	18.87 $\pm$	1.33 $\pm$ 0.03	3.56 $\pm$ 0.07	0.56 $\pm$ 0.02	0.34 $\pm$ 0.00	0.09 $\pm$ 0.01	0.13 $\pm$ 0.02
2	396 $\pm$ 7.6	0.84 $\pm$ 0.03	0.87 $\pm$ 0.01	263 $\pm$ 20	5.8 $\pm$ 0.5	8.42 $\pm$ 0.14	4.8 $\pm$ 0.04	0.76 $\pm$ 0.01	3.48 $\pm$ 0.00	0.08 $\pm$ 0.00	0.56 $\pm$ 0.00	0.23 $\pm$ 0.00	1.74 $\pm$ 0.03
3	334 $\pm$ 6.3	0.77 $\pm$ 0.03	0.89 $\pm$ 0.03	208 $\pm$ 8.3	13.2 $\pm$ 1.2	11.6 $\pm$ 0.07	10.2 $\pm$ 0.01	1.90 $\pm$ 0.35	2.76 $\pm$ 0.23	0.07 $\pm$ 0.00	0.20 $\pm$ 0.00	0.13 $\pm$ 0.00	0.06 $\pm$ 0.00
4	306 $\pm$ 4.1	0.78 $\pm$ 0.03	0.83 $\pm$ 0.02	181 $\pm$ 0.3	11.2 $\pm$ 0.7	17.2 $\pm$ 0.49	25.3 $\pm$ 0.49	1.27 $\pm$ 0.13	1.41 $\pm$ 0.01	0.19 $\pm$ 0.09	0.21 $\pm$ 0.05	0.25 $\pm$ 0.01	0.43 $\pm$ 0.01
5	300 $\pm$ 6.0	0.56 $\pm$ 0.03	0.63 $\pm$ 0.01	207 $\pm$ 2.8	11.5 $\pm$ 0.2	13.1 $\pm$ 0.07	9.3 $\pm$ 0.07	6.62 $\pm$ 0.01	2.93 $\pm$ 0.01	0.07 $\pm$ 0.02	0.16 $\pm$ 0.00	0.10 $\pm$ 0.01	0.09 $\pm$ 0.01
6	286 $\pm$ 6.5	0.61 $\pm$ 0.02	0.72 $\pm$ 0.02	256 $\pm$ 1.5	13.5 $\pm$ 0.0	24.0 $\pm$ 0.42	22.5 $\pm$ 0.92	1.03 $\pm$ 0.01	2.95 $\pm$ 0.00	0.08 $\pm$ 0.00	0.25 $\pm$ 0.00	0.19 $\pm$ 0.00	0.09 $\pm$ 0.00
7	278 $\pm$ 8.3	0.52 $\pm$ 0.03	0.57 $\pm$ 0.02	227 $\pm$ 7.8	13.3 $\pm$ 0.6	21.27 $\pm$ 1.13	11.1 $\pm$ 0.42	3.91 $\pm$ 0.05	3.18 $\pm$ 0.02	0.07 $\pm$ 0.00	0.19 $\pm$ 0.00	0.13 $\pm$ 0.00	0.06 $\pm$ 0.00
8	276 $\pm$ 4.0	0.62 $\pm$ 0.05	0.65 $\pm$ 0.03	58 $\pm$ 0.7	3.8 $\pm$ 0.1	8.0 $\pm$ 0.28	10.7 $\pm$ 0.18	1.36 $\pm$ 0.00	2.79 $\pm$ 0.03	0.07 $\pm$ 0.00	0.79 $\pm$ 0.02	0.2 $\pm$ 0.01	0.02 $\pm$ 0.00
9	269 $\pm$ 5.3	0.57 $\pm$ 0.01	0.59 $\pm$ 0.02	330 $\pm$ 7.9	22.4 $\pm$ 0.2	4.71 $\pm$ 0.08	3.7 $\pm$ 0.01	3.49 $\pm$ 0.01	5.23 $\pm$ 0.04	0.07 $\pm$ 0.00	0.79 $\pm$ 0.01	0.16 $\pm$ 0.00	0.39 $\pm$ 0.01
10	268 $\pm$ 7.5	0.49 $\pm$ 0.05	0.54 $\pm$ 0.02	217 $\pm$ 10	11.0 $\pm$ 0.6	27 $\pm$ 1.2	16.3 $\pm$ 0.57	10.8 $\pm$ 0.13	5.74 $\pm$ 0.10	0.14 $\pm$ 0.06	0.44 $\pm$ 0.02	0.21 $\pm$ 0.00	0.24 $\pm$ 0.01
11	265 $\pm$ 8.2	0.68 $\pm$ 0.03	0.79 $\pm$ 0.03	134 $\pm$ 6.6	5.7 $\pm$ 0.4	10.9 $\pm$ 0.14	18.2 $\pm$ 0.07	1.87 $\pm$ 0.01	3.97 $\pm$ 0.02	0.11 $\pm$ 0.01	0.36 $\pm$ 0.00	0.16 $\pm$ 0.00	0.07 $\pm$ 0.00
12	254 $\pm$ 6.8	0.60 $\pm$ 0.00	0.65 $\pm$ 0.01	172 $\pm$ 4.9	25.6 $\pm$ 0.7	8.1 $\pm$ 0.14	8.2 $\pm$ 0.07	1.71 $\pm$ 0.01	3.38 $\pm$ 0.02	0.07 $\pm$ 0.00	0.20 $\pm$ 0.00	0.14 $\pm$ 0.00	0.05 $\pm$ 0.00
13	237 $\pm$ 6.0	0.51 $\pm$ 0.02	0.59 $\pm$ 0.02	252 $\pm$ 9.5	11.9 $\pm$ 0.3	30.75 $\pm$ 0.99	16.6 $\pm$ 0.35	1.39 $\pm$ 0.01	3.6 $\pm$ 0.04	0.09 $\pm$ 0.00	0.34 $\pm$ 0.01	0.18 $\pm$ 0.00	0.19 $\pm$ 0.01
14	228 $\pm$ 5.6	0.50 $\pm$ 0.03	0.61 $\pm$ 0.03	209 $\pm$ 2.0	11.3 $\pm$ 0.1	17.7 $\pm$ 0.49	22.3 $\pm$ 0.49	1.46 $\pm$ 0.03	4.74 $\pm$ 0.07	0.07 $\pm$ 0.00	0.25 $\pm$ 0.00	0.23 $\pm$ 0.00	0.09 $\pm$ 0.01
15	224 $\pm$ 6.9	0.53 $\pm$ 0.01	0.61 $\pm$ 0.04	266 $\pm$ 10	7.3 $\pm$ 1.7	22.1 $\pm$ 0.3	25.7 $\pm$ 0.1	1.81 $\pm$ 0.02	4.6 $\pm$ 0.01	0.11 $\pm$ 0.02	0.16 $\pm$ 0.00	0.14 $\pm$ 0.00	0.12 $\pm$ 0.00
16	205 $\pm$ 5.9	0.48 $\pm$ 0.03	0.52 $\pm$ 0.02	182 $\pm$ 5.2	6.3 $\pm$ 0.5	11.1 $\pm$ 0.03	14.1 $\pm$ 0.7	1.2 $\pm$ 0.01	3.33 $\pm$ 0.06	0.09 $\pm$ 0.00	0.01 $\pm$ 0.02	0.03 $\pm$ 0.02	0.4 $\pm$ 0.09
17	205 $\pm$ 0.7	0.50 $\pm$ 0.00	0.40 $\pm$ 0.01	280 $\pm$ 4.7	21.2 $\pm$ 0.4	4.26 $\pm$ 0.30	4.1 $\pm$ 0.10	0.56 $\pm$ 0.06	6.62 $\pm$ 0.18	0.17 $\pm$ 0.08	0.39 $\pm$ 0.09	0.15 $\pm$ 0.01	0.01 $\pm$ 0.00
18	197 $\pm$ 12.4	0.42 $\pm$ 0.01	0.54 $\pm$ 0.02	232 $\pm$ 9.7	7.0 $\pm$ 0.4	15.5 $\pm$ 0.00	11.8 $\pm$ 0.01	0.53 $\pm$ 0.01	3.07 $\pm$ 0.01	0.19 $\pm$ 0.09	0.27 $\pm$ 0.00	0.12 $\pm$ 0.00	0.02 $\pm$ 0.00
19	195 $\pm$ 9.2	0.41 $\pm$ 0.02	0.46 $\pm$ 0.00	259 $\pm$ 13.6	7.8 $\pm$ 0.3	20.1 $\pm$ 0.21	17.2 $\pm$ 0.14	11.3 $\pm$ 0.04	4.35 $\pm$ 0.04	0.07 $\pm$ 0.00	0.14 $\pm$ 0.00	0.15 $\pm$ 0.00	0.03 $\pm$ 0.00
20	194 $\pm$ 8.7	0.35 $\pm$ 0.03	0.40 $\pm$ 0.01	256 $\pm$ 3.3	12.2 $\pm$ 0.6	29.3 $\pm$ 0.07	15.2 $\pm$ 0.00	3.51 $\pm$ 0.01	3.82 $\pm$ 0.00	0.09 $\pm$ 0.00	0.56 $\pm$ 0.01	0.24 $\pm$ 0.00	0.21 $\pm$ 0.00
21	191 $\pm$ 3.9	0.44 $\pm$ 0.01	0.48 $\pm$ 0.00	218 $\pm$ 0.6	6.6 $\pm$ 0.1	12.7 $\pm$ 0.14	8.9 $\pm$ 0.05	1.90 $\pm$ 0.00	3.62 $\pm$ 0.01	0.09 $\pm$ 0.00	0.41 $\pm$ 0.00	0.13 $\pm$ 0.00	0.06 $\pm$ 0.00
22	179 $\pm$ 2.7	0.41 $\pm$ 0.06	0.52 $\pm$ 0.03	199 $\pm$ 11.2	6.7 $\pm$ 0.1	16.7 $\pm$ 0.21	18.7 $\pm$ 0.21	2.14 $\pm$ 0.01	13.92 $\pm$ 0.06	0.08 $\pm$ 0.00	0.36 $\pm$ 0.00	0.18 $\pm$ 0.00	0.05 $\pm$ 0.00
23	177 $\pm$ 3.7	0.45 $\pm$ 0.02	0.52 $\pm$ 0.02	120 $\pm$ 3.4	10.5 $\pm$ 0.5	5.38 $\pm$ 0.19	6.6 $\pm$ 0.07	1.47 $\pm$ 0.01	7.51 $\pm$ 0.09	0.07 $\pm$ 0.00	0.17 $\pm$ 0.00	0.12 $\pm$ 0.00	n.d
24	166 $\pm$ 3.0	0.37 $\pm$ 0.03	0.43 $\pm$ 0.01	192 $\pm$ 21	6.3 $\pm$ 0.6	38.36 $\pm$ 0.49	25.4 $\pm$ 0.14	2.0 $\pm$ 0.01	3.89 $\pm$ 0.01	0.08 $\pm$ 0.00	0.37 $\pm$ 0.00	0.17 $\pm$ 0.00	0.06 $\pm$ 0.00
25	163 $\pm$ 6.7	0.35 $\pm$ 0.02	0.40 $\pm$ 0.01	256 $\pm$ 4.7	8.3 $\pm$ 0.5	7.38 $\pm$ 0.99	4.6 $\pm$ 0.29	1.75 $\pm$ 0.04	2.86 $\pm$ 0.14	0.08 $\pm$ 0.00	0.34 $\pm$ 0.16	0.11 $\pm$ 0.00	0.01 $\pm$ 0.00
26	162 $\pm$ 2.0	0.43 $\pm$ 0.02	0.46 $\pm$ 0.02	177 $\pm$ 1.8	4.4 $\pm$ 0.3	4.05 $\pm$ 0.54	5.6 $\pm$ 0.34	0.50 $\pm$ 0.00	5.34 $\pm$ 0.03	0.08 $\pm$ 0.01	0.34 $\pm$ 0.06	0.14 $\pm$ 0.00	0.42 $\pm$ 0.06
27	159 $\pm$ 12.6	0.40 $\pm$ 0.01	0.43 $\pm$ 0.03	173 $\pm$ 11	13 $\pm$ 0.7	9.7 $\pm$ 0.42	11.2 $\pm$ 0.24	1.78 $\pm$ 0.02	1.11 $\pm$ 0.02	0.08 $\pm$ 0.00	0.16 $\pm$ 0.00	0.14 $\pm$ 0.00	0.06 $\pm$ 0.00
28	157 $\pm$ 6.6	0.39 $\pm$ 0.01	0.44 $\pm$ 0.01	147 $\pm$ 6.6	13.5 $\pm$ 0.2	8.7 $\pm$ 0.42	8.2 $\pm$ 0.16	0.74 $\pm$ 0.00	2.77 $\pm$ 0.00	0.09 $\pm$ 0.01	0.31 $\pm$ 0.08	0.16 $\pm$ 0.00	0.17 $\pm$ 0.01
29	156 $\pm$ 9.0	0.40 $\pm$ 0.03	0.47 $\pm$ 0.02	133 $\pm$ 2.3	9.3 $\pm$ 0.2	19.3 $\pm$ 0.01	25.3 $\pm$ 0.14	0.65 $\pm$ 0.00	3,3 $\pm$ 0.00	0.09 $\pm$ 0.00	0.21 $\pm$ 0.00	0.14 $\pm$ 0.00	0.09 $\pm$ 0.00
30	156 $\pm$ 1.8	0.36 $\pm$ 0.01	0.40 $\pm$ 0.00	239 $\pm$ 6.4	9.5 $\pm$ 1.6	14.9 $\pm$ 0.78	13.3 $\pm$ 0.42	0.96 $\pm$ 0.01	3.79 $\pm$ 0.08	0.08 $\pm$ 0.00	0.37 $\pm$ 0.03	0.16 $\pm$ 0.00	0.08 $\pm$ 0.00
31	153 $\pm$ 3.5	0.33 $\pm$ 0.03	0.35 $\pm$ 0.01	215 $\pm$ 6.1	7.3 $\pm$ 0.2	13.5 $\pm$ 0.14	9.6 $\pm$ 0.04	0.76 $\pm$ 0.01	2.19 $\pm$ 0.01	0.08 $\pm$ 0.00	0.34 $\pm$ 0.00	0.16 $\pm$ 0.00	0.03 $\pm$ 0.00

Table 4.6 (continued): FRAP, TEAC, total phenolic, α -tocopherol, γ -tocopherol and phenolic contents of collected olive oil samples. Values are expressed as mean $\pm$ SD.

Sample	TP	TEAC	FRAP	α -Toc	γ -Toc	TY	HTY	OP	Pino	CA	VA	FA	p-COU
32	153 $\pm$ 9.9	0.41 $\pm$ 0.01	0.47 $\pm$ 0.01	160 $\pm$ 3.7	9.3 $\pm$ 0.6	9.19 $\pm$ 0.35	12.4 $\pm$ 0.28	0.52 $\pm$ 0.00	3.06 $\pm$ 0.06	0.08 $\pm$ 0.00	0.25 $\pm$ 0.00	0.13 $\pm$ 0.00	0.10 $\pm$ 0.01
33	147 $\pm$ 1.8	0.32 $\pm$ 0.04	0.39 $\pm$ 0.02	202 $\pm$ 6.8	7.9 $\pm$ 0.4	12.1 $\pm$ 0.28	6.3 $\pm$ 0.28	0.59 $\pm$ 0.01	3.85 $\pm$ 0.00	0.07 $\pm$ 0.00	0.61 $\pm$ 0.00	0.16 $\pm$ 0.00	0.08 $\pm$ 0.01
34	145 $\pm$ 3.4	0.38 $\pm$ 0.02	0.40 $\pm$ 0.01	193 $\pm$ 12	15.1 $\pm$ 0.3	10.9 $\pm$ 0.49	17.3 $\pm$ 0.42	0.59 $\pm$ 0.00	2.38 $\pm$ 0.04	0.08 $\pm$ 0.00	0.25 $\pm$ 0.00	0.17 $\pm$ 0.00	0.14 $\pm$ 0.01
35	145 $\pm$ 2.7	0.37 $\pm$ 0.02	0.42 $\pm$ 0.01	193 $\pm$ 4.9	14.5 $\pm$ 1.0	30.1 $\pm$ 0.92	19.3 $\pm$ 0.42	0.49 $\pm$ 0.01	2.56 $\pm$ 0.04	0.12 $\pm$ 0.00	0.30 $\pm$ 0.02	0.14 $\pm$ 0.00	0.12 $\pm$ 0.01
36	144 $\pm$ 2.5	0.34 $\pm$ 0.00	0.37 $\pm$ 0.01	164 $\pm$ 1.1	13.7 $\pm$ 0.5	10.8 $\pm$ 0.00	9.9 $\pm$ 0.03	1.99 $\pm$ 0.01	1.29 $\pm$ 0.02	0.08 $\pm$ 0.00	0.19 $\pm$ 0.00	0.16 $\pm$ 0.00	0.09 $\pm$ 0.00
37	138 $\pm$ 2.4	0.33 $\pm$ 0.03	0.42 $\pm$ 0.01	240 $\pm$ 27	15.7 $\pm$ 1.9	10.6 $\pm$ 0.28	12.3 $\pm$ 0.14	1.11 $\pm$ 0.01	10.78 $\pm$ 0.12	0.08 $\pm$ 0.00	0.39 $\pm$ 0.02	0.19 $\pm$ 0.00	0.30 $\pm$ 0.01
38	129 $\pm$ 5.8	0.28 $\pm$ 0.01	0.30 $\pm$ 0.01	154 $\pm$ 2.5	4.0 $\pm$ 1.1	3.11 $\pm$ 0.07	3.4 $\pm$ 0.04	0.51 $\pm$ 0.00	4.87 $\pm$ 0.03	0.07 $\pm$ 0.00	0.26 $\pm$ 0.01	0.14 $\pm$ 0.00	0.29 $\pm$ 0.01
39	128 $\pm$ 1.4	0.19 $\pm$ 0.02	0.24 $\pm$ 0.01	289 $\pm$ 9.3	16.3 $\pm$ 0.4	23.5 $\pm$ 0.21	3.7 $\pm$ 0.01	0.37 $\pm$ 0.00	4.15 $\pm$ 0.02	0.09 $\pm$ 0.00	1.45 $\pm$ 0.01	0.24 $\pm$ 0.00	0.13 $\pm$ 0.01
40	126 $\pm$ 11.1	0.28 $\pm$ 0.02	0.34 $\pm$ 0.01	120 $\pm$ 0.2	4.2 $\pm$ 0.3	19.8 $\pm$ 0.07	6.8 $\pm$ 0.01	2.09 $\pm$ 0.00	3.10 $\pm$ 0.01	0.07 $\pm$ 0.00	0.20 $\pm$ 0.00	0.14 $\pm$ 0.00	0.10 $\pm$ 0.00
41	124 $\pm$ 0.9	0.22 $\pm$ 0.02	0.29 $\pm$ 0.01	264 $\pm$ 17	6.3 $\pm$ 0.4	7.3 $\pm$ 0.14	3.9 $\pm$ 0.04	0.34 $\pm$ 0.02	3.5 $\pm$ 0.03	0.08 $\pm$ 0.00	0.48 $\pm$ 0.01	0.16 $\pm$ 0.00	0.07 $\pm$ 0.00
42	122 $\pm$ 3.3	0.33 $\pm$ 0.01	0.43 $\pm$ 0.01	209 $\pm$ 4.4	6.5 $\pm$ 0.2	3.96 $\pm$ 0.11	4.6 $\pm$ 0.10	0.41 $\pm$ 0.00	4.57 $\pm$ 0.01	0.09 $\pm$ 0.00	0.60 $\pm$ 0.00	0.18 $\pm$ 0.00	0.16 $\pm$ 0.00
43	121 $\pm$ 3.4	0.29 $\pm$ 0.01	0.34 $\pm$ 0.00	202 $\pm$ 0.3	14.7 $\pm$ 0.7	28.9 $\pm$ 1.06	3.0 $\pm$ 0.04	0.42 $\pm$ 0.00	2.27 $\pm$ 0.04	0.10 $\pm$ 0.00	0.23 $\pm$ 0.01	0.13 $\pm$ 0.00	0.11 $\pm$ 0.00
44	118 $\pm$ 2.8	0.15 $\pm$ 0.02	0.16 $\pm$ 0.00	97 $\pm$ 2.6	4.8 $\pm$ 0.1	20.4 $\pm$ 1.56	1.1 $\pm$ 0.01	0.68 $\pm$ 0.06	1.43 $\pm$ 0.04	0.17 $\pm$ 0.08	0.20 $\pm$ 0.02	0.15 $\pm$ 0.01	0.16 $\pm$ 0.01
45	116 $\pm$ 1.9	0.25 $\pm$ 0.00	0.30 $\pm$ 0.00	240 $\pm$ 9.3	8.0 $\pm$ 0.4	7.32 $\pm$ 0.14	3.5 $\pm$ 0.04	1.57 $\pm$ 0.02	3.11 $\pm$ 0.01	0.09 $\pm$ 0.01	0.20 $\pm$ 0.00	0.12 $\pm$ 0.00	n.d
46	114 $\pm$ 5.7	0.25 $\pm$ 0.01	0.32 $\pm$ 0.01	281 $\pm$ 8.5	12.0 $\pm$ 0.3	8.64 $\pm$ 0.35	3.6 $\pm$ 0.07	0.38 $\pm$ 0.01	2.94 $\pm$ 0.04	0.08 $\pm$ 0.00	0.20 $\pm$ 0.00	0.15 $\pm$ 0.00	0.50 $\pm$ 0.01
47	112 $\pm$ 2.3	0.26 $\pm$ 0.02	0.32 $\pm$ 0.01	171 $\pm$ 1.5	6.3 $\pm$ 0.2	8.2 $\pm$ 0.07	6.6 $\pm$ 0.09	0.45 $\pm$ 0.00	6.02 $\pm$ 0.00	0.08 $\pm$ 0.00	0.51 $\pm$ 0.00	0.20 $\pm$ 0.00	0.26 $\pm$ 0.00
48	109 $\pm$ 2.5	0.14 $\pm$ 0.02	0.22 $\pm$ 0.00	38 $\pm$ 0.2	4.0 $\pm$ 0.1	3.61 $\pm$ 0.05	0.2 $\pm$ 0.04	0.49 $\pm$ 0.00	4.32 $\pm$ 0.04	0.07 $\pm$ 0.00	0.26 $\pm$ 0.01	0.13 $\pm$ 0.00	n.d
49	107 $\pm$ 1.3	0.33 $\pm$ 0.03	0.24 $\pm$ 0.01	167 $\pm$ 6.7	9.3 $\pm$ 0.3	16.2 $\pm$ 0.49	6.7 $\pm$ 0.14	2.64 $\pm$ 0.55	2.17 $\pm$ 0.02	0.64 $\pm$ 0.09	0.30 $\pm$ 0.17	0.19 $\pm$ 0.08	0.42 $\pm$ 0.02
50	104 $\pm$ 3.8	0.27 $\pm$ 0.02	0.30 $\pm$ 0.01	152 $\pm$ 5.1	3.7 $\pm$ 0.2	7.1 $\pm$ 0.00	7.7 $\pm$ 0.01	2.7 $\pm$ 0.01	3.5 $\pm$ 0.01	0.08 $\pm$ 0.00	0.16 $\pm$ 0.00	0.14 $\pm$ 0.00	0.14 $\pm$ 0.01
51	101 $\pm$ 1.1	0.23 $\pm$ 0.00	0.27 $\pm$ 0.01	181 $\pm$ 7.0	8.0 $\pm$ 0.3	4.60 $\pm$ 0.04	3.5 $\pm$ 0.01	2.0 $\pm$ 0.00	1.87 $\pm$ 0.00	0.08 $\pm$ 0.01	0.16 $\pm$ 0.00	0.12 $\pm$ 0.00	0.01 $\pm$ 0.00
52	100 $\pm$ 0.4	0.27 $\pm$ 0.01	0.36 $\pm$ 0.00	164 $\pm$ 1.1	13.7 $\pm$ 0.5	4.1 $\pm$ 0.16	4.1 $\pm$ 0.04	0.38 $\pm$ 0.01	4.49 $\pm$ 0.06	0.08 $\pm$ 0.00	0.62 $\pm$ 0.01	0.19 $\pm$ 0.00	0.16 $\pm$ 0.01
53	91 $\pm$ 1.3	0.16 $\pm$ 0.01	0.21 $\pm$ 0.00	201 $\pm$ 4.5	5.9 $\pm$ 0.1	6.7 $\pm$ 0.57	2.8 $\pm$ 0.12	0.59 $\pm$ 0.04	2.17 $\pm$ 0.08	0.07 $\pm$ 0.00	0.17 $\pm$ 0.00	0.13 $\pm$ 0.00	0.04 $\pm$ 0.00
54	88 $\pm$ 0.9	0.18 $\pm$ 0.02	0.26 $\pm$ 0.00	138 $\pm$ 2.1	5.7 $\pm$ 1.2	7.9 $\pm$ 0.28	2.7 $\pm$ 0.05	0.42 $\pm$ 0.00	2.13 $\pm$ 0.04	0.07 $\pm$ 0.00	0.20 $\pm$ 0.02	0.17 $\pm$ 0.00	0.63 $\pm$ 0.03
55	83 $\pm$ 2.6	0.18 $\pm$ 0.01	0.22 $\pm$ 0.00	209 $\pm$ 1.0	14.8 $\pm$ 0.2	11.7 $\pm$ 0.42	3.0 $\pm$ 0.04	0.39 $\pm$ 0.01	1.50 $\pm$ 0.02	0.09 $\pm$ 0.00	0.46 $\pm$ 0.00	0.16 $\pm$ 0.01	0.30 $\pm$ 0.02
56	69 $\pm$ 1.1	0.18 $\pm$ 0.01	0.22 $\pm$ 0.01	188 $\pm$ 1.0	7.3 $\pm$ 0.1	4.51 $\pm$ 0.07	3.8 $\pm$ 0.03	0.49 $\pm$ 0.00	4.26 $\pm$ 0.02	0.07 $\pm$ 0.00	0.15 $\pm$ 0.00	0.12 $\pm$ 0.00	0.08 $\pm$ 0.00
57	45 $\pm$ 3.2	0.11 $\pm$ 0.01	0.18 $\pm$ 0.00	197 $\pm$ 15	13.2 $\pm$ 1.3	7.03 $\pm$ 0.05	0.54 $\pm$ 0.01	1.19 $\pm$ 0.03	1.86 $\pm$ 0.02	0.91 $\pm$ 0.03	0.15 $\pm$ 0.01	0.08 $\pm$ 0.02	0.19 $\pm$ 0.00

Abbreviations: TP; total phenolic, α -Toc; alpha tocopherol, γ -Toc; gamma tocopherol, TY; tyrosol, HTY; hydroxytyrosol, OP; oleuropein, Pino; pinoresinol, FA; ferulic acid, CA; caffeic acid, VA; vanillic acid, p-COU; p-coumaric acid, n.d; not detected. Values are expressed as mg/kg

4.3 Animal Study

In this study, we have focused on investigating the effects of olive oil naturally rich in phenolic compounds on several oxidative stress and inflammation parameters in SAMP8 mice. It is important to take into account that in the present study, we used a Western type diet with 20% fat, in which 10% fat replaced by olive oil and 0.15% cholesterol to trigger the oxidative stress in mice. Western type diet is a human diet of many countries associated with increased risk of atherosclerosis and CHD (Fung et al., 2009; Perona et al., 2010). The content of antioxidant vitamins was kept low but adequate and synthetic antioxidants and flavonoids were completely excluded from the diet in order to minimize their impact on the outcome of the experiment. The two oils that are used in the study were identical for major and minor constituents (Table 4.7).

Table 4.7: Concentrations of phenols and TEAC and FRAP values of the high-polyphenol (HP) and low-polyphenol (LP) olive oils.

	High-polyphenol olive oil	Low-polyphenol olive oil
Phenolic compounds [mg/kg oil]		
Tyrosol	20.78	7.03
Hydroxytyrosol	18.87	0.54
Pinoresinol	3.56	1.85
Oleuropein	1.33	1.19
Caffeic acid	0.56	0.91
Vanillic acid	0.34	0.16
p-coumaric acid	0.13	0.20
Ferulic acid	0.09	0.08
α -Tocopherol	217	197
Total phenolics [mg GA equivalent/kg oil]	532	44
FRAP [mmol AA equivalent/kg oil]	1.27	0.17
TEAC [mmol trolox equivalent/kg oil]	1.01	0.11

4.3.1 Body weight and feed intake

At the end of the 4.5-month feeding trial, no differences were observed in the final body weight and daily feed intake of mice fed the HP and LP diets, as shown in Table 4.8.

Table 4.8: Body weight and food intake of SAMR1 and SAMP8 mice fed either with high or low phenolic diet.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
Body Weight (g)	31.15±0.45	29.12±1.31	27.19±1.43
Food intake (g/day)	2.42±0.01	3.13±0.07	3.26±0.48

Values are expressed as mean±SEM (n=10)

4.3.2 Effect of age and HP diet on serum parameters

In serum samples of mice PON1 activity was measured as a marker of the age-associated oxidative status. Arylesterase activity was measured, which is more closely related to the levels of PON1 protein (Costa et al., 2005) thus, more indicative of the status of the PON1 enzyme. Significant decrease was found in aged SAMP8 LP mice as compared to SAMR1 LP mice ($p<0.001$). Feeding the mice with a HP diet increased the PON-1 activity in serum of mice ($p<0.05$, Figure 4.1). The change in PON-1 activity may stem from either HDL levels or PON1 transcription in liver. But no significant difference was found either in serum HDL level between groups (Table 4.9) or PON-1 mRNA expression in liver (Figure 4.7).

According to our results the difference in arylesterase activity cannot be attributed to changes in these parameters due to the fact that no significant difference of HDL levels in serum and PON-1 mRNA expression levels were determined in liver. This decline in PON-1 activity may be related to increased oxidative stress with aging (Seres et al., 2004) and the increase in PON-1 activity may be related to effect of HP diet.

In serum, total triglyceride levels did not differ among groups on the other hand total cholesterol levels were significantly decreased in SAMP8 LP mice ($p<0.001$). The effect of age and diet on serum biochemical parameters (total cholesterol, total triglyceride and HDL) is given in Table 4.9.




Figure 4.1: Mean ($\pm$ SEM) PON1 activity in serum of SAMR1 and SAMP8 mice fed a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either LP or HP (n=9).

Table 4.9 : Serum parameters of SAMR1 and SAMP8 mice.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
HDL [mg/dl]	92 $\pm$ 8.1	63.3 $\pm$ 15	59.5 $\pm$ 11
Total cholesterol [mg/dl]	160 $\pm$ 3.1	116 $\pm$ 6.9***	105 $\pm$ 2.9
Total triglyceride [mg/dl]	120 $\pm$ 13	142 $\pm$ 13	110 $\pm$ 9.3

Values are expressed as mean $\pm$ SEM (n=6-10) ***p<0.001 SAMR1-LP vs SAMP8-LP.

4.3.3 Effect of age and hp diet on oxidative stress biomarkers

In order to determine the effect of ageing on lipid peroxidation and protein oxidation, TBARS, protein carbonyl levels and proteasomal activity were determined in liver and heart tissues of mice. In liver significantly higher TBARS ($p<0.01$) and protein carbonyl levels ($p<0.05$) were obtained in SAMP8 LP mice as compared to SAMR1 LP mice (Table 4.10). No effect of HP diet was observed on TBARS levels but level of protein carbonyls decreased with HP treatment ($p<0.05$). In order to detect if the change in protein carbonyl level is related to proteasomal activity, we measured the activity of subunits of proteasomal system. Proteasomal activity of the β -1, β -2 and β -5 subunits in liver tissue remained unchanged (Table 4.10).

In heart TBARS levels were interestingly lower in SAMP8 LP mice as compared to SAMR1 mice ($p<0.05$). As observed in liver, treatment with phenolic rich olive oil significantly decreased TBARS levels of SAMP8 mice as compared to SAMP8 LP mice ($p<0.001$, Table). Protein carbonyl levels were higher in SAMP8 LP than that of SAMR1 LP ($p<0.05$, Table 4.11) and a significant decrease was observed in with

HP diet as compared to LP diet ($p < 0.01$, Table 4.11). In terms of proteasomal activities, no significant difference was observed in terms of β -2 and β -5 subunits among groups, whereas activity of β -1 subunit was decreased in SAMP8 LP mice as compared to SAMR1 LP mice ($p < 0.05$). No effect of HP treatment was observed in SAMP8 mice ($p > 0.05$, Table 4.11).

Table 4.10: Concentrations of TBARS and protein carbonyls as well as proteasomal activities in liver homogenates from SAMR1 and SAMP8 mice fed a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either LP or HP for 4.5 months.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
TBARS [nmol/g tissue]	442 $\pm$ 35	700 $\pm$ 78**	663 $\pm$ 48
Protein carbonyls [nmol/g protein]	775 $\pm$ 36	874 $\pm$ 36*	730 $\pm$ 46 ^{##}
Proteasomal activity [μ mol/mg*min]			
β -1 subunit	32.8 $\pm$ 2.1	31.1 $\pm$ 2.9	31.2 $\pm$ 6.7
β -2 subunit	6.2 $\pm$ 0.2	6.1 $\pm$ 0.1	5.9 $\pm$ 0.2
β -5 subunit	56.7 $\pm$ 5.7	40.9 $\pm$ 3.1	49.7 $\pm$ 4.2

Values are expressed as mean $\pm$ SEM (n=10) * $p < 0.05$ SAMR1-LP vs SAMP8-LP, ** $p < 0.01$ SAMR1-LP vs SAMP8-LP ^{##} $p < 0.01$ SAMP8-LP vs SAMP8-HP.

Table 4.11: Concentrations of TBARS and protein carbonyls as well as proteasomal activities in heart homogenates from SAMR1 and SAMP8 mice fed diets with 10% olive oil for 4.5 months.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
TBARS [nmol/g tissue]	563 $\pm$ 46	428 $\pm$ 27*	274 $\pm$ 22 ^{###}
Protein carbonyls [nmol/g protein]	763 $\pm$ 33	959 $\pm$ 70*	665 $\pm$ 37 ^{##}
Proteasomal activity [μ mol/mg*min]			
β -1 subunit	29.6 $\pm$ 1.0	26.1 $\pm$ 1.0*	31.7 $\pm$ 3.4
β -2 subunit	6.9 $\pm$ 0.4	8.2 $\pm$ 0.5	6.8 $\pm$ 0.5
β -5 subunit	72.2 $\pm$ 6.9	93.0 $\pm$ 9.2	79.3 $\pm$ 11

Values are expressed as mean $\pm$ SEM (n=10) * $p < 0.05$ SAMR1-LP vs SAMP8-LP, ^{##} $p < 0.01$ SAMP8-LP vs SAMP8-HP, ^{###} $p < 0.001$ SAMP8-LP vs SAMP8-HP.

Depending on the age and analyzed organs different profiles can be observed in literature regarding TBARS and protein carbonyl levels of SAMP8 mice. Nabeshi et al. (2006) determined higher protein carbonyl levels in liver at 3 and 6 month of age however results did not differ at 9 month of age, and in brain higher levels were detected at 3 month of age but no difference was found at 6 and 9 month of age. Higher TBARS level was observed in brain of SAMP8 mice as compared to ddY mice at 11 month of age but no difference was found at 3 month of age. Matsugo et al. (2000) found increased TBARS level in heart at 3 and 6 month of age but the

results did not differ in SAMP8 mice of 9 month of age. In the study performed by Okatani et al. (2002) no difference was observed in TBARS level of SAMP8 brain at the age of 6 and 12 month but protein carbonyl levels significantly increased. In another study no age related difference was found in terms of TBARS and protein carbonyls in brain of SAMP8 mice (Sato et al., 1996). Similarly no effect of treatment and age on protein carbonyl levels was found in the heart of SAMP8 mice by Schiborr et al. (2010).

As a general consensus the decline in proteasomal activity is one of the parameters for oxidative stress, and aging but controversy results are obtained depending on the age, tissue, and different region of tissues. In addition three catalytic sites of proteasome can be differentially affected in aged tissues. It has been reported that proteasomal activity decreased significantly with age in lung and liver samples but not in heart, kidney and skeletal muscle (Breusing et al., 2009). El mohsen et al. (2004) reported that proteasomal enzyme activity was not affected by age in any of the brain regions (cortex, cerebellum and hippocampus) of rats. Also no difference between subunits were found in young and aged Wistar rats (Gavilan et al., 2009), Fisher 344 rats (Kavazis et al., 2007). Age-dependent inhibition of β -5 subunit activity in the retinal pigment epithelium of Fischer 344 rats were reported (Kapphahn et al., 2007). Lower expression of β -2 and β -5 catalytic subunits in peripheral blood T lymphocytes was detected during aging (Ponnapannan et al., 2007). Activities of β -1 and β -2 subunits were diminished with age, while β -5 activity did not exhibit significant alterations in different age groups of Fisher 344 rat hearts (Bulteau et al., 2002). Age-dependent protein modifications and declining proteasome activity in the human lens detected in all catalytic subunits of age group >60 but just β -1 subunit activity declined in age group of 25-55 years (Viteri et al., 2004). By contrast there was a 3-fold increase (Ferrington et al., 2005) and 4- fold increase in β -1 and β -5 subunits in aged muscle of Fischer 344 rats (Husom et al., 2004).

4.3.4 Effect of age and HP diet on redox status

As indicators of the redox status, we measured concentrations of the GSH and oxidized GSH (GSSG), amino thiols (methionine, homocysteine and cysteine) and the antioxidant uric acid in liver and heart. Homocysteine level could not be measured in both tissues as the concentration was above the standard range. No difference was found in GSH, GSSG, GSH/GSSG, methionine and uric acid between groups in liver of mice (Table 4.12). Cysteine level was significantly higher in SAMP8 LP as compared to SAMR1 group ($p < 0.05$).

Table 4.12: The concentration of GSH, GSSG, GSH/GSSG, cysteine, methionine and uric acid in liver of SAMR1 and SAMP8 mice at 7 month of age.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
GSH [nmol/mg protein]	28.9±5.2	31.9±3.0	38.6±5.9
GSSG [nmol/mg protein]	0.63±0.11	0.94±0.14	1.05±0.13
GSH/GSSG [nmol/mg protein]	38.1±3.7	31.3±4.0	37.2±3.2
Cysteine [nmol/mg protein]	0.50±0.07	0.68±0.05*	0.75±0.08
Methionine [nmol/mg protein]	1.68±0.14	1.61±0.15	1.74±0.36
Uric Acid [nmol/mg protein]	0.67±0.13	0.88±0.10	0.68±0.09

Values are expressed as mean± SEM (n=10). *p<0.05 SAMR1-LP vs SAMP8-LP.

In heart GSSG and thus GSH/GSSG could not be measured as the concentration was below the calibration range. No difference was found in GSH and methionine level between groups in heart of mice (Table 4.13). Cysteine and uric acid levels were significantly higher in SAMP8 LP as compared to SAMR1 group (p<0.05). Feeding the mice with HP diet decreased both parameters significantly (p<0.05) as compared to SAMP8 LP mice.

Table 4.13: The concentration of GSH, GSSG, GSH/GSSG, cysteine, methionine and uric acid in heart of SAMR1 and SAMP8 mice at 7 month of age.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
GSH [nmol/mg protein]	3.97±0.57	4.80±0.57	3.66±0.74
Cysteine [nmol/mg protein]	0.14±0.03	0.22±0.04*	0.14±0.03 [#]
Methionine [nmol/mg protein]	0.27±0.04	0.39±0.08	0.20±0.06
Uric Acid [nmol/mg protein]	0.25±0.04	0.55±0.13*	0.23±0.04 [#]

Values are expressed as mean± SEM (n=10). No significant difference was found.

Cysteine is a sulfhydryl-containing amino acid which shows cytotoxicity in vitro and promotes detachment of human arterial endothelial cells in culture (Dudman et al., 1991). It also exhibits autooxidation properties in the presence of metal ions resulting in the generation of free radicals and hydrogen peroxide (Hogg, 1999). It is indicated that cysteine could be atherogenic as it supports superoxide-mediated modification of LDL, thus facilitating foam cell formation (Verhoef et al., 1996; El-Khairi et al., 2001). Furthermore increased cysteine levels associate with cardiovascular diseases (Mills et al., 2000; Jacob et al., 1999). In addition to its antioxidant activity, uric acid is another cardiovascular risk factor which can promote LDL oxidation in vitro and liberate peroxide and superoxide free radical. Hyperuricemia is considered to be a risk factor in many clinical conditions including hypertension, cardiovascular diseases (Nakanishi et al., 2003; Strazzullo and Puig,

2007). In our study elevated levels of cysteine and uric acid was detected in heart of SAMP8 LP group as compared to SAMR1 LP group indicating a susceptibility of SAMP8 for cardiovascular diseases ($p < 0.05$). Phenolic rich diet improved the both parameters significantly ($p < 0.05$, Table 4.13). Schiborr et al. (2010) did not find any significant difference either between the hearts of SAMP8 and SAMR1 mice or treatment groups regarding cysteine and uric acid levels. Rebrin and Sohal (2004) reported significantly lower levels of cysteine in blood of SAMP mice as compared to C57BL/6 mice at 13 month of age.

4.3.5 Effect of age and HP diet on antioxidant vitamins in liver

Age related alterations of two important endogenous vitamins, ascorbic acid and α -tocopherol, which are related to inflammation and oxidative stress were also measured. Ascorbic acid is an essential micronutrient in primates which exhibits some of its biological activity due to its free radical scavenging activity. It is a water soluble chain breaking radical scavenger and recycles plasma membrane α -tocopherol via the reduction of the α -tocopheroxyl radical (May, 2000). It exhibited a 2 fold increase in SAMP8 LP mice than those in SAMR1 LP. No change was observed in vitamin C level in heart of SAMP8 and SAMR1 mice (Figure 4.2 A-B).




Figure 4.2 A-B: Mean ($\pm$ SEM) vitamin C concentration (nmol/mg protein) in liver A) in heart B) of SAMR1 and SAMP8 mice fed a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or high amount of phenolics (n=10).

The mechanisms for increased ascorbic acid levels may include increased synthesis and/or transport of that vitamin (Siqueira et al., 2011). Ascorbic acid is transported into mammalian cells by two types of proteins: the sodium-ascorbate co-transporters, SVCT1 and SVCT2, which cause Na^+ -dependent uptake of L-ascorbate. Gluconolactone oxidase (Gulo) is the last enzyme in ascorbic acid synthesis. Therefore the mRNA expression level of these genes was also measured. The level of SVCT1 mRNA expression was significantly increased in SAMP8 LP mice as compared to SAMR1 but no change was determined in terms of SVCT2 and Gulo (Table). Treatment with HP diet increased Gulo mRNA expression as compared to SAMP8 LP mice but SVCT1 and SVCT2 remained unchanged (Table 4.14).

Table 4.14: Relative mRNA expression (normalized for GAPDH) of SVCT1, SVCT2 and Gulonolactone oxidase in liver of SAMR1 and SAMP8 mice fed for 4.5 months a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or high amount of phenolics. Mice were killed at 7 months of age.

	SAMR1 LP	SAMP8 LP	SAMP8 HP
SVCT1	0.62±0.05	0.84±0.09*	0.80±0.08
SVCT2	0.70±0.14	0.55±0.10	0.70±0.09
Gulonolactone oxidase	0.54±0.07	0.59±0.08	0.88±0.07 [#]

Values are expressed as mean ± SEM (n=10). *p<0.05 SAMR1-LP vs SAMP8-LP, [#]p<0.05 SAMP8-LP vs SAMP8-HP.

As ascorbic acid recycles α -tocopherol, we determined the age- and ascorbic acid related change of that lipid soluble vitamin. In the same manner α -tocopherol exhibited a 2 fold increase in SAMP8 LP mice than those in SAMR1 LP mice (Figure 4.3). Also mice fed HP diet exhibited increased α -tocopherol levels as compared to SAMP8 LP mice (Figure 4.3).




Figure 4.3: Mean (±SEM) vitamin E concentration (mmol/mg protein) in liver of SAMR1 and SAMP8 mice fed a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or high amount of phenolics (n=10).

As far as the antioxidant vitamins concerned ascorbic acid content significantly increased in SAMP8 LP mice. We measured the mRNA level of genes related to ascorbic acid transport or synthesis as an eventual mechanism of the increase. The mRNA expression of dominant ascorbic acid transporter in liver (SVCT1) increased in SAMP8 LP mice. However the expression of SVCT2 remained unchanged in aged mice. This result is in accordance with the study of Michels et al (2003) who

found no age related change in rat hepatocytes. It has been reported that primary role of SVCT1 is maintenance of the whole-body homeostasis, through dietary absorption and renal reabsorption, while SVCT2 is crucial for ascorbic acid uptake in metabolically active and specialized tissues, such as eye, brain, skeletal muscle and heart (Savini et al., 2008). Beside Kuo et al. (2004) showed that SVCTs appear to function independently of each other, as SVCT1 expression and the ascorbate concentration in SVCT1-predominant organs such as liver, was not affected by SVCT2 deficiency. Furthermore no difference was found in Gulo levels in our study suggesting that this increase may not be attributed to de novo ascorbic acid synthesis. Also Lykkesfeldt et al. (1998) reported no age associated alteration in rat hepatic ascorbic acid synthesis.

In addition increased level of α -tocopherol was also observed in our aged SAMP8 mice. This result is in agreement with other studies (Battisti et al., 1994; van der Loo et al., 2002; Petusottir et al., 2007). It is suggested that both ascorbic acid and α -tocopherol levels of SAMP8 mice may increase to counteract the negative effects of age related oxidative stress with their antioxidant activities. The increase in α -tocopherol levels in HP group could not be explained with increased vitamin C level. It is supposed that this increase could be due to difference in vitamin E metabolites.

The membrane bound tocoperoxyl radical may be reduced back to tocopherol by ascorbate (Vitamin C). In turn, the ascorbyl radical may be regenerated to ascorbate by thiol or polyphenol antioxidants within the so-called antioxidant network (Packer et al., 2001). Besides the increase in tocopherol levels may be due to the genes that take part in vitamin E metabolism. Therefore we measured the mRNA expression levels of genes in vitamin E metabolism. As shown in Figure 4.4 A-H no significant change was found for all the genes analysed except the expression of MDR-1 in SAMR1 LP and SAMP8 LP groups, which means the increase in vitamin E is may be due to the regeneration effect of vitamin C.




Figure 4.4 A-H: Relative mRNA expression of genes in vitamin E metabolism (normalized for GAPDH) of CYP3A (A), TTP (B), LPL (C), LRP1 (D), SRB1 (E), MDR-1 (F), LDLR (G), ABCA1(H) in liver of SAMP8 and SAMR1 mice.

Present data suggest that SAMP8 mice increase hepatic ascorbic acid uptake via SVCT1 as an adaptive response mechanisms. Increased tissue levels of ascorbic acid may in turn affect Nrf2 dependent gene expression thereby impairing stress response via enzymatic antioxidant defence mechanisms. Ristow et al. (2010) have recently demonstrated that ascorbic acid may interfere with physical exercise related induction of the Nrf2 target genes superoxide dismutase and glutathione peroxidase in humans.

Apart from their antioxidant properties, prooxidant activities of ascorbic acid and α -tocopherol have been described. Ascorbic acid promotes the Fenton reaction in vitro and may accelerate non enzymatic lipid peroxidation in various tissues (Halliwell, 1996). It is also reported that ascorbic acid increases inflammation in ascorbic acid deficient mice (Ki et al., 2010), down regulate key genes connected with free radical scavenging and repair (Selman et al., 2006) and dose dependently causes exacerbating stress contributing to liver injury (Abdul-Rezzak et al., 2010). α -tocopherol gives rise to the formation of tocopheroxyl radicals in increased oxidative stress conditions (Rietjens et al., 2002), it can catalyze hydroperoxide-induced hepatocyte lipid peroxidation and cytotoxicity (Tafazoli et al., 2005). Alternatively high ascorbic acid and α -tocopherol content in SAMP8 mice may also facilitate inflammation and oxidative stress acting as a prooxidant in our study. Further studies are required to measure ascorbic acid and α -tocopherol levels in other tissues of SAMP8 mice to see whether our findings in liver are in consistent with the other tissues regarding oxidative damage. Also the amount of tocopheroxyl radicals can be measured in liver to support our statement.

4.3.6 Effect of age and HP diet on antioxidant phase II enzymes

Increased attention has been given to induction of phase II enzymes as an important mechanism for chemoprevention against carcinogenesis, but the studies in terms of correlation between antioxidant enzymes and aging are scarce.

In the present study, we examined expression levels of γ -GCS, GST- α , NQO-1, HO-1, PON-1 (in liver) and PON-2 (in heart) genes as Phase II enzymes in heart and liver to evaluate the antioxidant capacity. These genes are modulated by ARE (antioxidant response element) via binding of Nrf2 transcription factor. Nrf2 is an important molecular switch, which orchestrates the gene expression of antioxidant and phase II enzymes (Lewis et al., 2010; Sykiotis et al., 2011). Little is known about the role of olive oil phenolics on Nrf2 dependent gene expression in vivo.

In our study no significant difference was found between SAMR1 LP and SAMP8 LP mice in terms of γ -GCS, GST- α , PON-2 and Nrf2 expression levels. On the contrary NQO-1 mRNA level was significantly lower ($p<0.001$) and HO-1 level was significantly higher ($p<0.05$) in SAMP8 LP group. Treatment with phenolic rich olive oil improved the levels of γ -GCS and GST- α ($p<0.01$), NQO-1, PON-2 and Nrf2 ($p<0.05$) significantly in heart (Figure 4.5). No effect was observed in HO-1 level.




Figure 4.5 A-F: Relative mRNA expression (normalized for GAPDH) of Nrf2 (A), GST, (B), γ -GCS, (C) NQO-1, (D), HO-1 (E), and PON-2 (F) in hearts of SAMP8 mice fed for 4.5 months a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or high amount of phenolics. Values are expressed as mean $\pm$ SEM (n=10).

In liver no difference was detected between mice groups regarding Nrf2 modulated genes. Only HO-1 mRNA expression was significantly lower ($p<0.05$) in SAMP8 LP group as compared to SAMR1 LP group (Figure 4.7). This decrease was also shown in protein level by Western blot (Figure 4.6). We detected a tendency for γ -GCS and GST- α genes in mRNA level. Therefore we detected the protein level of both genes but similarly as observed in mRNA level, no difference was found for the protein levels of γ -GCS and GST- α (Figure 4.8).




Figure 4.6: Protein levels of HO-1. Protein levels were determined via western blotting. Shown is a representative western blot. Values are expressed as the mean $\pm$ SEM (n=10) ** $p<0.01$ SAMR1 vs SAMP8.




Figure 4.7 A-F: Relative mRNA expression (normalized for GAPDH) of Nrf2 (A), GST, (B), γ-GCS, (C), NQO- 1, (D), HO-1 (E), and PON-2 (F) in livers of SAMP8 mice fed for 4.5 months Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or high amount of phenolics. Mice were killed at 7 months of age. Values are expressed as mean ± SEM (n=10).




Figure 4.8 A-B: Protein levels of GST-α (A) and γ-GCS (B). Protein levels were determined via western blotting. Shown is a representative western blot. Values are expressed as the mean $\pm$ SEM (n=10).

Protective effect of Nrf2 mediated antioxidant enzyme activities on cardiovascular diseases and vascular health has been documented (Zhu et al., 2005; Mann et al., 2007; Mann et al., 2009; Hur et al., 2010). It is suggested that olive oil phenolics may exert their modulatory effects on aging, cardiovascular health through Nrf2/ARE pathway. It has been recently shown that hydroxytyrosol induces antioxidant/detoxicant enzymes and Nrf2 translocation in hepatocytes (Martin et al., 2010). Furthermore hydroxytyrosol induced cytoprotection against oxidative injury in vascular endothelial cells via Nrf2 dependent signal transduction pathway (Zrelli et al., 2011).

In studies regarding Nrf2-related genes that were conducted with aged rodents, no modulation of Nrf2 was observed neither with mulberry nor bilberry extract in SAMP8 mice (Shih et al, 2009). Suh et al. (2004) showed reduced Nrf2/ARE binding and Nrf2 mediated gene expression and NQO-1 levels in aging rat liver.

4.3.7 Effect of age and HP diet on SIRT1 expression

SIRT1, is a class III deacetylase, that can modulate numerous longevity-promoting cellular functions, endothelial function, lipid metabolism, cell cycling, gene expression, cardioprotection and aging (Longo and Kennedy, 2006). Previously decreased level of SIRT1 expression was documented in SAMP8 mice as compared

to its control SAMR1 mice (Pallas et al., 2008; Gutierrez-Cuesta et al., 2008; Tajés et al., 2010). In accordance with these studies, the SIRT1 expression level was significantly decreased in SAMP8 LP group ($p < 0.001$, Figure 4.9) but high phenolic olive oil treatment significantly increased mRNA expression levels in SAMP8 mice in our study ($p < 0.05$). SIRT1 exerts its functions by deacetylating variety of metabolic transcription factors. A moderate induction of SIRT retarded ageing of the heart and induced resistance to oxidative stress (Alcendor et al., 2007). Thus, the decreased concentrations of biomarkers of oxidative stress in heart of mice fed the HP diet might be related to the moderate SIRT1 induction observed in these animals.




Figure 4.9 A-B: Relative mRNA expression (normalized for GAPDH) of SIRT1 in liver (A) and heart (B) of mice fed for 4.5 months a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or and high amount of phenolics. Mice were killed at 7 months of age. Values are expressed as mean $\pm$ SEM (n=10).

4.3.8 Effect of age and HP diet on inflammation markers

As aging is accompanied by an increase in circulating acute phase reactants we measured the CRP and SAP mRNA expression levels as a marker of inflammatory status and an increased risk of cardiovascular diseases. Increased levels of CRP and SAP together with decreased level of HO-1 indicate that SAMP8 mice have higher inflammatory state. These results may also affect PON1 activity, which reduced during acute phase response (Ayub et al., 1999). Dullaart et al. (2009) reported a lower PON-1 activity related to higher plasma CRP levels in type 2

diabetic patients. As shown in figure 4.10 both mRNA expression levels of CRP and SAP increased in SAMP8 LP mice but no effect of HP treatment was observed.




Figure 4.10: Relative mRNA expression (normalized for GAPDH) of SAP and CRP levels in livers of SAMP8 and SAMR1 mice fed for 4.5 months a Western type diet with 0.15% cholesterol and 20% fat, in which 10% of fat was from olive oil containing either low or high amount of phenolics. Mice were killed at 7 months of age. Values are expressed as mean $\pm$ SEM (n=10).

5. CONCLUSION

The aim of the present study was to investigate the effect of a diet rich in phenolic compounds on the ageing related changes in human model of ageing, SAMP8 mice. Most of the studies have focused on the brain ageing of SAMP8 mice and effective results are obtained but studies regarding the heart and liver ageing of this mice are scarce and needs to be studied. The liver and heart are the major organs responsible for a number of physiological processes that are crucial to sustaining life. Another aim was to develop and validate a method for the quantification of olive oil phenolics with sensitive electrochemical detector.

As a result the effect of ageing was remarkable in liver of SAMP8 mice as compared to SAMR1 mice. This study demonstrates that age dependent oxidative stress and inflammation are elevated in SAMP8 mice from a quite early age as compared to its control SAMR1 mice which may lead to a rise in free radical formation and thus age related diseases. The increase in the protein and lipid oxidation may contribute to age related changes as reflected in PON1 activity and expression of inflammation related genes (CRP, SAP, HO-1). The amount of vitamin E and vitamin C was also higher in SAMP8 LP mice. On the other hand redox status, Nrf2 and Nrf2-dependent gene expressions remained to be unchanged in liver. The effect of HP diet was not observed except in TBARS and protein carbonyl levels.

In heart the effect of HP diet was obvious in SAMP8 HP. Feeding a HP diet reduced biomarkers of oxidative stress, induced Nrf2 and Nrf2- dependent gene expression and increased SIRT1 gene expression in the heart of SAMP8 mice compared to animals fed a LP diet. In the literature the effect of Nrf2 dependent antioxidant enzyme activities on cardiovascular diseases and vascular health is well known. Overall, our results suggest that a diet rich in olive oil phenolics reduces oxidative stress in the heart of SAMP8 mice most likely by induction of Nrf2-dependent gene expression. Mice fed the HP diet had lower concentrations of TBARS and protein carbonyls in the heart than mice on the LP diet. However, proteasomal activity was not changed. Thus differences in heart protein carbonyl levels were independent of proteasomal activity. Regarding Nrf2 and Nrf2-dependent gene expression and redox status no effect was detected in SAMP8 LP mice as compared to SAMR1 LP mice.

The HP and LP olive oils used in this study had similar profile but they notably differed in their content of the major olive oil phenolics hydroxytyrosol, tyrosol, oleuropein and pinoresinol which were reflected in the corresponding FRAP and TEAC values of the two oils. Importantly, the HP olive oil contained 12-fold higher concentrations of total phenolics than the LP olive oil. As far as the main olive oil phenolic compound, hydroxytyrosol, is considered, HP oil contains approximately 40 fold higher hydroxytyrosol than the LP oil. It is suggested that the observed beneficial effects of HP diet may be due to high hydroxytyrosol content, which seems to be of importance for the potential health benefits of olive oils.

In conclusion differences in tissue responses to the oxidative stress, different profiles of investigated biomarkers were obtained in heart and liver tissue of the mice. Some tissues respond to the environmental changes quickly but some tissues needs more stressed conditions. We fed the mice with a diet containing 20% fat and 0.15% cholesterol in order to trigger the oxidative stress. The 9-10 weeks old female mice fed either HP or LP diet for 4.5 months and were killed at 7 months old. In the literature different results have been obtained depending on the age of SAMP8 mice. Besides the results differ depending on the experimental design, type of the bioactive compound and the amount of the compounds. According to the results of our study, it is suggested that the mice were not old enough to see clear effects of phenolic rich diet therefore they should be fed longer period of time or older mice should be used.

Another factor for that may be the daily consumption of phenolic compounds. The amount of total phenolic compound in HP oil is 532 mg/kg and the diet contains 10% olive oil. Thus the amount of phenolics in the diet is 53.2 mg/kg. If the daily feed intake of the mice is taken into account, ~2 grams, which means daily amount of phenolic consumption is 0.1 mg per mouse. It is a quite low amount to show the clear beneficial effects of olive oil phenolics. The differences in tissues of mice fed either LP or HP diet may be attributed to low amount of phenolic compound consumption.

In further studies it is suggested to use different ageing models to study the effect of olive oil phenolics on ageing and life span. Besides different tissues of SAMP8 mice should be considered in order to understand the pathological processes in ageing of other tissues.

In terms of our validated method it is possible to analyze selected phenolic compounds (tyrosol, hydroxytyrosol, oleuropein, pinoresinol, caffeic acid, vanillic

acid, p-coumaric acid, ferulic acid) in 17 minutes. This method was validated in terms of selectivity, lower limit of quantification and detection, accuracy, precision, recovery, short term stability, freeze thaw stability and post preparative stability. The phenolics were quantified in ng level. The recoveries of these phenolic compounds ranged between 72-96%. This method may also be used in biological samples (urine, blood, tissues) to analyse the phenolics and as well as their metabolites.

REFERENCES

- Aaby, K., Skrede, G. and Wrolstad, R.E.** 2005. Phenolic composition and antioxidant activities in flesh and achenes of strawberries (*Fragaria ananassa*). *Journal of Agricultural and Food Chemistry*, **53**, 4032-4040.
- Abe, R., Beckett, J., Abe, R., Nixon, A., Rochier, A., Yamashita, N. and Sumpio, B.** 2011. Olive oil polyphenol oleuropein inhibits smooth muscle cell proliferation. *European Journal of Vascular and Endovascular* **41**, 814-820.
- Abdul-Razzak, K.K., Alzoubi, K.H., Abdo, S.A. and Hananeh, W.M.** 2010. High-dose vitamin C: Does it exacerbate the effect of psychosocial stress on liver? Biochemical and histological study. *Experimental Toxicology and Pathology*, doi: 10.1016/j.etp.2010.09.011
- Achilli, G., Cellerino, G.P. and Hamache, P.H.** 1993. Identification and determination of phenolic constituents in natural beverages and plant extracts by means of a coulometric electrode array system. *Journal of Chromatography*, **632**, 111-117.
- Al-Azzawie, H.F. and Alhamdani, M.S.S.** 2006. Hypoglycemic and antioxidant effect of oleuropein in alloxan-diabetic rabbit. *Life Science* **78**, 1371-1377.
- Alcendor, R.R., Gao, S., Zhai, P., Zablocki, D., Holle, E., Yu, X., Tian, B., Wagner, T., Vatner, S.F. and Sadoshima, J.** 2007. Sirt1 regulates aging and resistance to oxidative stress in the heart. *Circulation Research*, **100**, 1512-1521
- Allameh, A., Maleklou, N., Zargar, M., and Sanati, M.H.** 2008. The influence of uric acid treatments on liver glutathione system prevent oxidative damages in experimental autoimmune encephalomyelitis mice *Neuroscience Letters*, **439**, 111-115.
- Álvarez-García, O., Vega-Naredo, I., Sierra, V., Caballero, B., Tomás-Zapico, C., Camins, A., García, J.J., Palls, M. and Coto-Montes, A.** 2006. Elevated oxidative stress in the brain of senescence-accelerated mice at 5 months of age. *Biogerontology*, **7**, 43-52.
- Alonso, A. and Martínez-González, M.A.** 2004. Olive oil consumption and reduced incidence of hypertension: the SUN study. *Lipids*, **39**, 1233-1238.
- Andreadou, I., Sigala, F., Iliodromitis, E.K., Papaefthimiou, M., Sigalas, C. and Aligiannis, N.** 2007. Acute doxorubicin cardiotoxicity is successfully treated with the phytochemical oleuropein through suppression of oxidative and nitrosative stress. *Journal of Molecular Cell and Cardiology*, **42**, 549-558.
- Andreadou, I., Iliodromitis, E.K., Mikros, E., Constantinou, M., Agalias, A. and Magiatis, P.** 2006. The olive constituent oleuropein exhibits anti-ischemic, antioxidative, and hypolipidemic effects in anesthetized rabbits. *Journal of Nutrition*, **136**, 2213-2219.

- Angerosa, F., d'Alessandro, N., Konstantinou, P. and Di Giacinto, L. 1995. GC-MS Evaluation of phenolic compounds in virgin olive oil. *Journal of Agricultural and Food Chemistry*, **43**, 1802-1807.
- Aoyama, K., Matsumura, N., Watabe, M., Wang, F., Kikuchi-Utsumi, K. and Nakaki, T. 2011. Caffeine and uric acid mediate glutathione synthesis for neuroprotection. *Neuroscience*, **181**, 206-215.
- Aparicio, R. and Luna, R. 2002. Characterisation of monovarietal virgin olive oils. *European Journal of Lipid Science and Technology*, **104**, 614-627.
- Aviram, M., Rosenblat, M., Bisgaier, C.L., Newton, R.S., Primo-Parmo, S.L. and La Du, B.N. 1998. Paraoxonase inhibits high density lipoprotein oxidation and preserves its function: a possible peroxidative role for paraoxonase. *Journal of Clinical Investigations*, **101**, 1581-1590.
- Ayton, J., Mailer, R.J., Haigh, A., Tronson, D. and Conlan, D. 2007. Quality and oxidative stability of Australian olive oil according to harvest date and irrigation. *Journal of Food Lipids*, **14**, 138-156.
- Ayub, A., Mackness, M.I., Arrol, S., Mackness, B., Patel, J. and Durrington, P.N., 1999. Serum paraoxonase after myocardial infarction. *Arteriosclerosis Thrombosis Vascular Biology*, **19**, 330-335.
- Banning, A., Deubel, S., Kluth, D., Zhou, Z. and Brigelius-Flohe, R. 2005. The GI-GPx gene is a target for Nrf2. *Molecular Cell Biology*, **25**, 4914-4923.
- Barberger-Gateau, P., Raffaitin, C., Letenneur, L., Berr, C., Tzourio, C. and Dartigues, J.F. 2007. Dietary patterns and risk of dementia: the Three-City cohort study. *Neurology*, **69**, 1921-1930.
- Barja, G., 2004. Free radicals and aging. *Trends in Neuroscience*, **27**, 595-600.
- Battisti, C., Dotti, M.T., Manneschi, L. and Federico, A. 1994. Increase of serum levels of vitamin E during human aging: is it a protective factor against death? *Archives of Gerontology and Geriatrics*, suppl. **4**, 13-18.
- Bazoti, F.N., Bergquist, J., Markides, K. and Tsarbopoulos, A. 2008. Localization of the noncovalent binding site between amyloid- β -peptide and oleuropein using electrospray ionization FT-ICR mass spectrometry *Journal of American Society of Mass Spectrometry*, **19**, 1078-1085.
- Becker, B.F. 1993. Towards the physiological function of uric acid. *Free Radical Biology and Medicine* **14**, 615-631.
- Bello, R.I., Gómez-Díaz, C., Burón, M.I., Navas, P. and Villalba, J.M. 2006. Differential regulation of hepatic apoptotic pathways by dietary olive and sunflower oils in the aging rat. *Experimental Gerontology*, **41**, 1174-1184.
- Beltran, G., Aguilera, M.P., Del Rio, C., Sanchez, S. and Martinez, L. 2005. Influence of fruit ripening process on the natural antioxidant content of Hojiblanca virgin olive oils. *Food Chemistry*, **89**, 207-215.
- Bendini, A., Bonoli, M., Cerretani, L., Biguzzi, B., Lercker, G. and Toschi, T.G. 2003. Liquid-liquid and solid-phase extractions of phenols from virgin olive oil and their separation by chromatographic and electrophoretic methods. *Journal of Chromatography A*, **985**, 425-433.

- Benzie, I.F. and Strain, J.J.** 1996. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": The FRAP assay. *Analytical Biochemistry*, **239**, 70-76
- Bernstein, P.S.** 2002. New insights into the role of the macular carotenoids in age-related macular degeneration. Resonance Raman studies. *Pure Applied Chemistry*, **74**, 1419-1425.
- Bertelli, A.A., Migliori, M., Panichi, V., Longoni, B., Origlia, N. and Ferretti, A.** 2002. Oxidative stress and inflammatory reaction modulation by white wine. *Annals of the New York Academy of Sciences*, **957**, 295-301.
- Bianco, A., Coccioli, F., Guiso, M. and Marra, C.** 2001. The occurrence in olive oil of a new class of phenolic compounds: hydroxy-isochromas. *Food Chemistry*, **77**, 450.
- Blumberg J.** 2004. Use of biomarkers of oxidative stress in research studies. *Journal of Nutrition*, **134**, 3188-3189.
- Bogani, P., Galli, C., Villa, M. and Visioli, F.** 2007. Postprandial anti-inflammatory and antioxidant effects of extra virgin olive oil. *Atherosclerosis*, **190**, 181-186.
- Boldyrev, A., Yuneva, M., Sorokina, E., Kramarenko, G., Fedorova, T., Konovalova, G. and Lankin, V.** 2001. Antioxidant systems in tissues of Senescence Accelerated Mice (SAM) characterized by accelerated ageing, *Biochemistry (Moscow)*, **66**, 1157-1163.
- Bonoli, M., Bendini, A., Cerretani, L., Lercker, G. and Toschi, T.** 2004. Qualitative and semiquantitative analysis of phenolic compounds in extra virgin olive oils as a function of the ripening degree of olive fruits by different analytical techniques. *Journal of Agricultural and Food Chemistry*, **52**, 7026-7032.
- Bonora, E., Targher, G., Zenere, MB, Saggiani, F, Cacciatori, V. and Tosi, F.** 1996. Relationship of uric acid concentration to cardiovascular risk factors in young men. role of obesity and central fat distribution. the verona young men atherosclerosis risk factors study. *International Journal of Obesity and Related Metabolic Disorders*, **20**, 975-980.
- Boskou, D.** 2006. Olive Oil Composition, in *Olive Oil: Chemistry and Technology*, **4**, Eds, Boskou, D., AOCS Press, Champaign, IL, USA.
- Braga, C., La Vecchia, C., Franceschi, S.** 1998. Olive oil, other seasoning fats, and the risk of colorectal carcinoma. *Cancer*, **82**, 448-453.
- Breusing, N., Grune, T.** 2008. Regulation of proteasome-mediated protein degradation during oxidative stress and aging. *Biological Chemistry*, **389**, 203-209.
- Breusing, N., Arndt, J., Voss, P., Bresgen, N., Wiswedel, I., Gardemann, A., Siems, W. and Grune, T.** 2009. Inverse correlation of protein oxidation and proteasome activity in liver and lung. *Mechanisms of Ageing and Development*, **130**, 748-753
- Bokov, A., Chaudhuri, A. and Richardson, A.** 2004. The role of oxidative damage and stress in aging. *Mechanisms of Ageing and Development*, **125**, 811-826.
- Brenes, M., Garcia, A., Garcia, P. and Garrido, A.** 2000. Rapid and complete extraction of phenols from olive oil and determination by means of a

- coulometric electrode array system. *Journal of Agricultural and Food Chemistry*, **48**, 5178-5183.
- Briante, R., Patumi, M., Limongelli, S., Febbraio, F., Vaccaio, C., DiSalle, A., LaCara, F. and Nucci, R.** 2002. Changes in phenolic and enzymatic activities content during fruit ripening in two Italian cultivars of *Olea europaea*. *Plant Science*, **162**, 791-798.
- Brod, S.A.**, 2000. Unregulated inflammation shortens human functional longevity. *Inflammation Research*, **49**, 561–570.
- Bruunsgaard, H.**, 2006. The clinical impact of systemic low-level inflammation in elderly populations. With special reference to cardiovascular disease, dementia and mortality. *Danish Medical Bulletin*, **53**, 285–309.
- Bu, Y., Rho, S., Kim, J., Kim, M.Y., Lee, D.H., Kim, S.Y., Choi, H. and Kim, H.** 2007. Neuroprotective effect of tyrosol on transient focal cerebral ischemia in rats. *Neuroscience Letters*, **414**, 218–221.
- Bulteau, A.L., Szveda, L.I. and Friguet, B.** 2002. Age-Dependent Declines in Proteasome Activity in the Heart. *Archives of Biochemistry and Biophysics*, **397**, 298-304.
- Burton, G.W., Joyce, A. and Ingold, K. U.** 1982. First proof that vitamin E is major lipid-soluble, chain-breaking antioxidant in human blood plasma. *Lancet*, **293**, 327.
- Buss, H., Chan, T. P., Sluis, K. B., Domigan, N. M. and Winterbourn, C.C.** 1997. Protein carbonyl measurement by a sensitive ELISA method. *Free Radical Biology and Medicine*, **23**, 361-366.
- Butterfield, D.A. and Poon, H.F.** 2005. The senescence-accelerated prone mouse (SAMP8): A model of age-related cognitive decline with relevance to alterations of the gene expression and protein abnormalities in Alzheimer's disease. *Experimental Gerontology* **40**, 774-783.
- Carluccio, M.A., Siculella, L., Ancora, M.A., Massaro, M., Scoditti, E., Storelli, C., Visioli, F., Distanti, A. and de Caterina, R.** 2003. Olive oil and red wine antioxidant polyphenols inhibit endothelial activation: antiatherogenic properties of mediterranean diet phytochemicals. *Arteriosclerosis, Thrombosis and Vascular Biology*, **23**, 622-629.
- Carretero, M., Escames, G., López, L.C., Venegas, C., Dayoub, J.C., García, L. and Acuña-Castroviejo, D.** 2009. Long-term melatonin administration protects brain mitochondria from aging. *Journal of Pineal Research*, **47**, 192-200.
- Carrasco-Pancorbo, A., Cerretani, L., Bendini, A., Segura-Carretero, A., Gallina-Toschi, T. and Fernandez-Gutiérrez, A.** 2005. Analytical determination of polyphenols in olive oils. *Journal of Separation Science*, **28**, 837-858.
- Caruso, D., Berra, B., Giavarini, F., Cortesi, N., Fedeli, E. and Galli, G.** 1999. Effect of virgin olive oil phenolic compounds on in vitro oxidation of human low density lipoproteins. *Nutrition Metabolism and Cardiovascular Diseases*, **9**, 102–107.
- Casal, S., Malheiro, R., Sendas, A., Oliveira, B.P.P. and Pereira, J.A.,** 2010. Olive oil stability under deep-frying conditions. *Food and Chemical Toxicology*, **48**, 2972-2979.

- Chiba, Y., Shimada, A., Kumagai, N., Yoshikawa, K., Ishii, S., Furukawa, A., Takei, S., Sakura, M., Kawamura, N. and Hosokawa, M.** 2009. The senescence-accelerated mouse (SAM): a higher oxidative stress and age-dependent degenerative diseases model. *Neurochemistry Research*, **34**, 679-687.
- Choi, A.M.K.** 2001. Heme-oxygenase-1 protects the heart. *Circulation Research*, **89**, 105–109.
- Choi, H.K. and Ford, E.S.** 2007. Prevalence of the metabolic syndrome in individuals with hyperuricemia. *American Journal of Medicine*, **120**, 442-447.
- Christophoridou, S., Dais, P., Tseng, L.H. and Spraul, M.** 2005. Separation and identification of phenolic compounds in olive oil by coupling high-performance liquid chromatography with postcolumn solid-phase extraction to nuclear magnetic resonance spectroscopy (LC-SPE-NMR). *Journal of Agricultural and Food Chemistry*, **53**, 4667-4679.
- Chung, H.Y., Cesari, M., Anton, S., Marzetti, E., Giovannini, S., Seo, A.Y., Carter, C., Yu, B.P. and Leeuwenburgh, C.** 2009. Molecular inflammation: Underpinnings of aging and age-related diseases. *Ageing Research Reviews*, **8**, 18-30.
- Chung, S., Yao, H., Caito, S., Hwang, J.W., Arunachalam, G. and Rahman, I.** 2010. Regulation of SIRT1 in cellular functions: role of polyphenols. *Archives of Biochemistry and Biophysics*, **501**, 79-90.
- Cicerale, S., Conlan, X.A., Sinclair, A.J. and Keast, R.S.** 2009. Chemistry and health of olive oil phenolics. *Critical Reviews in Food Science and Nutrition*, **49**, 218-236.
- Clodoveo, M.L., Delcuratolo, D., Gomes, T. and Colelli, G.** 2007. Effect of different temperatures and storage atmospheres on Coratina olive oil quality. *Food Chemistry*, **102**, 571–576.
- Coni, E., Di Benedetto, R., Di Pasquale, M., Masella, R. Modesti, D. and Mattei, R.** 2000. Protective effect of oleuropein, an olive oil biophenol, on low density lipoprotein oxidizability in rabbits. *Lipids*, **35**, 45-54.
- Covas, M.I., de la Torre, K., Farré-Albaladejo, M., Kaikkonen, J., Fitó, M., López-Sabater, C., Pujadas-Bastardes, M.A., Joglar, J., Weinbrenner, T., Lamuela-Raventós, R.M. and de la Torre.** 2006. Postprandial LDL phenolic content and LDL oxidation are modulated by olive oil phenolic compounds in humans. *Free Radical Biology and Medicine*, **40**, 608-616.
- Covas, M.I.** 2009. Olive oil and cardiovascular health. *Journal of Cardiovascular Pharmacology*, **54**, 477-482.
- Criado, M.N., Morelló, J.R., Motilva, M.J. and Romero, M.P.** 2004. Effect of growing area on pigment and phenolic fractions of virgin olive oils of the arbequina variety in Spain. *Journal of American Oil Chemist's Society*, **81**, 633–640.
- Cuesta, S., Kireev, R., Forman, K., García, C., Escames, G., Ariznavarreta, C., Vara, E. and Tresguerres, J.A.F.** 2010. Melatonin improves inflammation processes in liver of senescence-accelerated prone male mice (SAMP8). *Experimental Gerontology*, **45**, 950-956.

- Dalle-Donne I, Rossi R, Colombo R, Giustarini D, and Milzani A. 2006. Biomarkers of oxidative damage in human disease. *Clinical Chemistry*, **52**, 601-623.
- Davies MJ, Fu S, Wang H, and Dean RT. 1999. Stable markers of oxidant damage to proteins and their application in the study of human disease. *Free Radical Biology and Medicine*, **27**, 1151-1163.
- Dehghan, A., van Hoek, M., Sijbrands, E.J., Hofman, A. and Witteman, J.C.M. 2008. High serum uric acid as a novel risk factor for type 2 diabetes. *Diabetes Care*, **31**, 361-362.
- Deiana, M., Corona, G., Incani, A., Loru, D., Rosa, A., Atzeri, A., Paola, M and Dessì, M. 2010. Protective effect of simple phenols from extra virgin olive oil against lipid peroxidation in intestinal Caco-2 cells. *Food and Chemical Toxicology*, **48**, 3008-3016.
- de la Puerta, R., Ruiz Gutierrez, V. and Hout, J.R. 1999. Inhibition of leukocyte 5-lipoxygenase by phenolics from virgin olive oil. *Biochemical Pharmacology*, **57**, 445-449.
- de la Puerta, R., Martinez Dominguez, M.E., Ruiz-Gutierrez, V., Flavill, J.A. and Hout, J.R. 2001. Effects of virgin olive oil phenolics on scavenging of reactive nitrogen species and upon nitrgic neurotransmission. *Life Sciences*, **69**, 1213-1222.
- de la Torre-Carbot, K., Chávez-Servín, J.L., Jaregui, O., Castellote, A.I., Lamuela-Raventós, R.M., Nurmi, T., Poulsen, H.E., Gaddi, A.V., Kaikkonen, J., Zunft, J.F., Kieseewetter, H., Fitó, M., Covas, M.I. and López-Sabater. M.C. 2010. Elevated circulating LDL phenol levels in men who consumed virgin rather than refined olive oil are associated with less oxidation of plasma LDL. *Journal of Nutrition*, **140**, 501-508.
- De Stefani E, Brennan P, Boffetta P, Ronco AL, Mendilaharsu M and Deneo-Pellegrini H. 2000. Vegetables, fruits, related dietary antioxidants, and risk of squamous cell carcinoma of the esophagus: a case-control study in Uruguay. *Nutrition and Cancer*, **38**, 23-29.
- Della Ragione, F., Cucciolla, V., Borriello, A., Della Pietra, V., Pontoni, G., Racioppi, L., Manna, C., Galletti, P. and Zappia, V. 2000. Hydroxytyrosol, a natural molecule occurring in olive oil, induces cytochrome c-dependent apoptosis. *Biochemical and Biophysical Research Communications*, **278**, 733-739.
- Devaraj, S., Autret, B.C. and Jialal, I. 2006. Reduced-calorie orange juice beverage with plant sterols lowers C-reactive protein concentrations and improves the lipid profile in human volunteers. *American Journal of Clinical Nutrition*, **84**, 756-761.
- Devarajan, A., Bourquard, N., Hama, S., Navab, M., Grijalva, V.R., Morvardi, S., Clarke, C. F., Vergnes, L., Reue, K., Teiber, J.F. and Reddy, S.T. 2011 Paraoxonase 2 deficiency alters mitochondrial function and exacerbates the development of atherosclerosis. *Antioxidants and Redox Signaling*, **14**, 341-351.
- Di Benedetto, R., Vari, R., Scazzocchio, B., Filesì, C., Santangelo, C., and Giovannini, C. 2007. Tyrosol, the major extra virgin olive oil compound, restored intracellular antioxidant defences in spite of its weak antioxidative effectiveness. *Nutritional Metabolism and Cardiovascular Diseases*, **17**, 535-545.

- O'Dowd, Y., Driss, F., My-Chan Dang, P., Elbim, C., Gougerot-Pocidalo, M.A., Pasquier, C. and El-Benn, J. 2004. Antioxidant effect of hydroxytyrosol, a polyphenol from olive oil: scavenging of hydrogen peroxide but not superoxide anion produced by human neutrophils. *Biochemical Pharmacology*, **68**, 2003-2008.
- Dullaart, R.P.F., de Vries, R., Sluiter, W.J. and Voorbij, H.A.M. 2009. High plasma C-reactive protein (CRP) is related to low paraoxonase-I (PON-I) activity independently of high leptin and low adiponectin in type 2 diabetes mellitus. *Clinical Endocrinology*, **70**, 221–226.
- El Mohsen, M.A., Marks, J., Kuhnle, G., Rice-Evans, C., Moore, K., Gibson, G., Debnam, E. and Srai, S.K. 2004. The differential tissue distribution of the citrus flavanone naringenin following gastric instillation. *Free Radical Research*, **38**, 1329-1340.
- El-Khairi, L., Ueland, P.M., Refsum, H., Graham, I.M. and Vollset, S.E. 2001. Plasma total cysteine as a risk factor for vascular disease: The European Concerted Action Project. *Circulation*, **103**, 2544-2549.
- Esposito, K., Marfella, R. and Ciotola, M. 2004. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA*, **292**, 1440-1446.
- Esti, M., Cinquanta, L. and La Notte, E. 1998. Phenolic compounds in different olive varieties. *Journal of Agricultural and Food Chemistry*, **46**, 32-35.
- Estruch, R., Martínez-González, M.A. and Corella, D. 2006. Effects of a Mediterranean style diet on cardiovascular risk factors: a randomized trial. *Annals of Internal Medicine*, **145**, 1-11
- Farooqui, T. and Farooqui, A.A. 2010. Aging: An important factor for the pathogenesis of neurodegenerative diseases *Mechanisms of Ageing and Development*, **130**, 203–215.
- Farr, S.A., Poon, H.F., Dogrukol-Ak, D., Drake, J., Banks, W.A., Eyerman, E., Butterfield, A. and Morley, J.E. 2003. The antioxidants α -lipoic acid and N-acetylcysteine reverse memory impairment and brain oxidative stress in aged SAMP8 mice. *Journal of Neurochemistry*, **84**, 1173-1183.
- Fenton, M., Huang, H.L., Hong, Y., Hawe, E., Kurz, D.J. and Erusalimsky, J.D. 2004. Early atherogenesis in senescence-accelerated mice. *Experimental Gerontology*, **39**, 115–122.
- Ferrari, A.U., Radaelli, A. and Centola, M. 2003. Aging and the cardiovascular system. *Journal of Applied Physiology*, **95**, 2591-2597
- Ferrington, D.A., Husom, A.D. and Thompson, L.V. 2005. Altered proteasome structure, function, and oxidation in aged muscle. *The FASEB Journal*, **19**, 644-646.
- Ferris, C.D., Jaffrey, S.R. and Sawa, A. 1999. Haem oxygenase-1 prevents cell death by regulating cellular iron. *Nature Cell Biology*, **1**, 152–157.
- Ferruzzi, M.G., Bohm, V., Courtney, P.D. and Schwartz, S.J. 2002. Antioxidant and antimutagenic activity of dietary chlorophyll derivatives determined by radical scavenging and bacterial reverse mutagenesis assays. *Journal of Food Science*, **67**, 2589-2595.

- Finkel, T. and Holbrook, N.J.** 2000. Oxidants, oxidative stress and the biology of ageing. *Nature*, **408**, 239-247.
- Fitó, M., Cladellas, M., de la Torre, R., Martí, J., Alcántara, M., Pujadas-Bastardes, M., Marrugat, J., Bruguera, J., López-Sabater, M.C., Vila, J. and Covas, M.I.** 2005. Antioxidant effect of virgin olive oil in patients with stable coronary heart disease: a randomized, crossover, controlled, clinical trial. *Atherosclerosis*, **181**, 149-158.
- Forman, K., Vara E., Garcia, C., Ariznavarreta, C., Escames, G. and Tresguerres, J.A.F.** 2010. Cardiological aging in SAM model: effect of chronic treatment with growth hormone. *Biogerontology*, **11**, 275-286.
- Fortunato, G., Di Taranto MD, Bracale UM, Del Guercio, L., Carbone, F., Mazzaccara, C., Morgante, A., D'Armiento, F.P. and Porcellini, M.** 2008. Decreased paraoxonase-2 expression in human carotids during the progression of atherosclerosis. *Arteriosclerosis Thrombosis and Vascular Biology*, **28**, 594-600.
- Fuhrman, B., Volkova, N. and Aviram, A.** 2006. Postprandial serum triacylglycerols and oxidative stress in mice after consumption of fish oil, soy oil or olive oil: Possible role for paraoxonase-1 triacylglycerol lipase-like activit. *Nutrition*, **22**, 922-930.
- Fung, T.T., Rexrode, K.M., Mantzoros, C.S., Manson, J.E., Willett, W.C. and Hu, F.B.** 2009. Mediterranean diet and incidence and mortality of coronary heart disease and stroke in women. *Circulation*, **119**, 1093-1100.
- García, J.J., Piñol-Ripol, G., Martínez-Ballarín, E., Fuentes-Broto, L., Miana-Mena, F.J., Venegas, C., Caballero, B., Escames, G., Coto-Montes, A. and Acuña-Castroviejo D.** 2010. Melatonin reduces membrane rigidity and oxidative damage in the brain of SAMP8 mice. *Neurobiology of Aging*, **32**, 2045-2054.
- García-González, D.L., Aparicio-Ruiz, R. and Aparicio, R.** 2008. Virgin olive oil-Chemical implications on quality and health. *European Journal of Lipid Science and Technology*, **110**, 602-607.
- Gavilán, M.P., Castaño, A., Torres, M., Portavella, M., Caballero, C., Jiménez, S., García-Martínez, A., Parrado, J., Vitorica, J. and Ruano, D.** 2009. Age-related increase in the immunoproteasome content in rat hippocampus: molecular and functional aspects. *Journal of Neurochemistry*, **108**, 260-272.
- Gill, C.I.R., Boyd, A., McDermott, E., McCann, M., Servili, M., Selvaggini, R., Taticchi, A., Esposto, S., Montedoro, G., McGlynn, H. and Rowland, I.** 2005. Potential anti-cancer effects of virgin olive oil phenols on colorectal carcinogenesis models in vitro. *International Journal of Cancer*, **117**, 1-7.
- Giovannini, C., Straface, E., Modesti, D., Coni, E., Cantafora, A. and De Vincenzi, M.** 1999. Tyrosol, the major olive oil biophenol, protects against oxidized-LDL-induced injury in Caco-2 cells. *Journal of Nutrition*, **129**, 1269-1277.
- Giovannini, L., Migliori, M., Filippi, C., Origlia, N., Panichi, V. and Falchi, M.** 2002. Inhibitory activity of the white wine compounds, tyrosol and caffeic acid, on lipopolysaccharideinduced tumor necrosis factor-

alpha release in human peripheral blood mononuclear cells. *International Journal of Tissue Reactions*, **24**, 53–56.

- Gomez-Alonso, S., Salvador, M.D. and Fregapane, G.** 2002. Phenolic compounds profile of cornicabra virgin olive oil. *Journal of Agricultural and Food Chemistry*, **50**, 6812-6817.
- Gomez-Rico, A., Salvador, M.D., Moriana, A., Perez, D., Olmedilla, N., Ribas, F. and Fregapane, G.** 2007. Influence of different irrigation strategies in a traditional Cornicabra cv. Olive orchard on virgin olive oil composition and quality. *Food Chemistry*, **100**, 568-578.
- Gong, Y., Liu, L., Xie, B., Liao, Y., Yang, E. and Sun, Z.** 2008. Ameliorative effects of lotus seedpod proanthocyanidins on cognitive deficits and oxidative damage in senescence-accelerated mice. *Behavioural Brain Research*, **194**, 100-107.
- González-Santiago, M., Martín-Bautista, E., Carrero, J.J., Fonollá, J., Baró, L., Bartolomé, M.V., Gil-Loyzaga, P. and López-Huertas, E.** 2006. One-month administration of hydroxytyrosol, a phenolic antioxidant present in olive oil, to hyperlipemic rabbits improves blood lipid profile, antioxidant status and reduces atherosclerosis development *Atherosclerosis*, **188**, 35-42.
- Gordon, M.H., Paiva-Martins, F. and Almeida, M.** 2001. Antioxidant activity of hydroxytyrosol acetate compared with that of other olive oil polyphenols. *Journal of Agricultural and Food Chemistry*, **49**, 2480–2485.
- Gutierrez, F., Jimenez, B., Ruiz, A. and Albi, M.A.** 1999. Effect of olive ripeness on the oxidative stability of virgin olive oil extracted from the varieties picual and hojiblanca and on the different components involved. *Journal of Agricultural and Food Chemistry*, **47**, 121-127.
- Gutierrez-Cuesta, J., Tajés, M., Jiménez, A., Coto-Montes, A., Camins, A. and Pallás, M.** 2008. Evaluation of potential pro-survival pathways regulated by melatonin in a murine senescence model. *Journal of Pineal Research*, **45**, 497-505.
- Gutierrez-Rosales, F., Rios, J.J. and Gomez-Rey, M.L.** 2003. Main polyphenols in the bitter taste of virgin olive oil. structural confirmation by on-line high-performance liquid chromatography electrospray ionization mass spectrometry. *Journal of Agricultural and Food Chemistry*, **51**, 6021-6025.
- Hagen, T.M.** 2003. Oxidative stress, redox imbalance, and the aging process. *Antioxidants and Redox Signaling*, **5**, 503-506.
- Hallikainen, M., Lyyra-Laitinen, T. and Laitinen, T.** 2006. Endothelial function in hypercholesterolemic subjects: effects of plant stanol and sterol esters. *Atherosclerosis*, **188**, 425-432.
- Halliwell, B.** 1996. Vitamin C: antioxidant or pro-oxidant in vivo? *Free Radical Research*, **25**, 439-454.
- Hao, J., Shen, W., Yu, G., Jiaa, H., Li, X., Feng, Z., Wang, Y., Weber, P., Wertz, K., Sharman, E. and Liu, J.** 2010. Hydroxytyrosol promotes mitochondrial biogenesis and mitochondrial function in 3T3-L1 adipocytes. *Journal of Nutritional Biochemistry*, **21**, 634-644.
- Harman, D.,** 2003. The free radical theory of aging. *Antioxidants and Redox Signalling*, **5**, 557–561.

- Hogg, N.** 1999. The effect of cyst(e)ine on the autooxidation of homocysteine. *Free Radical Biology Medicine*, **27**, 28-33.
- Huang, D., Ou, B. and Prior, R. L.** 2005. The chemistry behind antioxidant capacity assays. *Journal of Agricultural and Food Chemistry*, **53**, 1841-1856.
- Huebbe, P., Jofre-Monseny, L. and Rimbach, G.** 2009. Alpha-tocopherol transport in the lung is affected by the apoE Genotype-studies in transgenic apoE3 and apoE4 mice. *IUBMB Life*, **61**, 453-456.
- Hur, K.Y., Kim, S. H., Choi, M.A., Williams, D.R., Lee, Y.H., Kan, S.W., Yadav, U.C.S, Srivastav, S.K., Jung, M., Cho, J.W., Kim, S.G., Kan, E.S., Lee, E.J. and Lee, H.C.** 2010. Protective effects of magnesium lithospermate B against diabetic atherosclerosis via Nrf2-ARE-NQO1 transcriptional pathway. *Atherosclerosis*, **211**, 69-76.
- Idriss, N.K., Blann, A.D. and Lip, G.Y.H.** 2008. Hemoxygenase-1 in cardiovascular disease. *Journal of the American College of Cardiology*, **52**, 971-978
- Ismail, A.S., Stavroulakis, G. and Metzidakis, J.** 1999. Effect of irrigation on the quality characteristics of organic olive oil. *Acta Horticulture*, **474**, 687-690.
- Jacob, N., Bruckert, E., Giral, P., Foglietti, M. and Turpin, G.** 1999. Cysteine is a cardiovascular risk factor in hyperlipidemic patients. *Atherosclerosis*, **146**, 53-59.
- Jacomelli, M., Pitozzi, V., Zaid, M., Larrosa, M., Tonini, G., Martini, A., Urbani, S., Taicchi, A., Servili, M., Dolara, P. and Giovannelli, L.** 2010. Dietary extra-virgin olive oil rich in phenolic antioxidants and the aging process: long-term effects in the rat. *Journal of Nutritional Biochemistry*, **21**, 290-296.
- Jaiswal, A.K.** 2004. Nrf2 Signaling in coordinated activation of antioxidant gene expression. *Free Radical Biology and Medicine*, **36**, 1199-1207.
- Jaouad, L., C. de Guise, H. Berrougui, M. Cloutier, M. Isabelle, T. Fulop, H. Payette, and A. Khalil.** 2006. Age-related decrease in high density lipoproteins antioxidant activity is due to an alteration in the PON1's free sulfhydryl groups. *Atherosclerosis*. **185**, 191-200.
- Jódar, L., Mercken, E.M., Ariza, J., Younts, C., González-Reyes, J.A., Alcaín, F.J., Burón, I., de Cabo, R. and Villalba, J.M.** 2011. Genetic deletion of Nrf2 promotes immortalization and decreases lifespan of murine embryonic fibroblasts. *The Journals of Gerontology A: Biological Sciences*, **66**, 247-256.
- Johnson, R.J., Perez-Pozo, S.E. and Sautin, Y.Y.** 2009. Hypothesis: could excessive fructose intake and uric acid cause type 2 diabetes? *Endocrinology Reviews*, **30**, 96-116.
- Johnson, F.B., Sinclair, D.A. and Guarente, L.** 1999. Molecular biology of aging. *Cell*, **96**, 291-302.
- Kamat, J.P., Bloor, K.K. and Devasagayam, T.P.A.** 2000. Chlorophyllin as an effective antioxidant against membrane damage in vitro and ex vivo. *Biochimica et Biophysica Acta- Molecular and Cell Biology of Lipids*, **1487**:113-127.
- Kapfahn, R.J., Bigelow, E.J. and Ferrington, D.A.** 2007. Age-dependent inhibition of proteasome chymotrypsin-like activity in the retina. *Experimental Eye Research*, **84**, 646-654.

- Kalua, C.M., Bedgood, D.R., Bishop, A.G. and Prenzler, P.D. 2006. Changes in volatile and phenolic compounds with malaxation time and temperature during virgin olive oil production. *Journal of Agricultural and Food Chemistry*, **54**, 7641-7651.
- Kavazis, A.N., DeRuisseau, K.C., McClung, J.M., Whidden, M.A., Falk, D.J., Smuder, A.J., Sugiura, T. and Powers, S.K. 2007. Diaphragmatic proteasome function is maintained in the ageing Fisher344 rat. *Experimental Physiology*, **92**, 895–901.
- Kayden, H.J. and Traber, M.G. 1993. Absorption, lipoprotein transport, and regulation of plasma concentrations of vitamin E in humans. *Journal of Lipid Research*, **34**, 343–358.
- Ki, M.R., Lee, H.R., Park, J.K., Hong, H., Han, S.Y., You, S.Y., Lee, E.M., Kim, A.Y., Lee, S.S., Jeong, K.S. 2010. Ascorbate promotes carbon tetrachloride-induced hepatic injury in senescence marker protein 30-deficient mice by enhancing inflammation. *Journal of Nutritional Biochemistry*, **22**, 535-542.
- Kim, Y., Lian, F., Yeum, K.J., Chongviriyaphan, N., Choi, S.W., Russell, R.M. and Wang, X.D. 2007. The effects of combined antioxidant (beta-carotene, a-tocopherol and ascorbic acid) supplementation on antioxidant capacity, DNA single-strand breaks and levels of insulin-like growth factor-1/IGF-binding protein 3 in the ferret model of lung cancer. *International Journal of Cancer*, **120**, 1847-1854
- Kirkwood, T.B.L. and Austad, S.N. 2000. Why do we age? *Nature*. **408**, 233-238.
- Kishido, T., Unno, K., Yoshida, H., Choba, D., Fukutomi, R., Asahina, S., Iguchi, K., Oku, N. and Hoshino, M. 2007. Decline in glutathione peroxidase activity is a reason for brain senescence: consumption of green tea catechin prevents the decline in its activity and protein oxidative damage in ageing mouse brain *Biogerontology*, **8**, 423-430.
- Kline, K., Lawson, K.A., Yu, W.P. and Sanders, B.G. 2003. Vitamin E and breast cancer prevention: Current status and future potential. *Journal of Mammary Gland Biology*, **8**, 91-102.
- Komatsu, T., Chiba, T., Yamaza, H., Yamashita, K., Shimada, A., Hoshiyama, Y., Henm, T., Ohtani, H., Higami, Y., de Cabo, R., Ingram, D.K. and Shimokawa, I. 2008. Manipulation of caloric content but not diet composition, attenuates the deficit in learning and memory of senescence-accelerated mouse strain P8. *Experimental Gerontology*, **43**, 339-346.
- Kontogianni, M.D., Panagiotakos, D.B., Chrysoshoou, C., Pitsavos, C., Zampelas, A. and Stefanadis, C. 2007. The impact of olive oil consumption pattern on the risk of acute coronary syndromes: the CARDIO2000 case-control study. *Clinical Cardiology*, **30**, 125-129.
- Kregel, K.C. and Zhang, H.J. 2007. An integrated view of oxidative stress in aging: basic mechanisms, functional effects, and pathological considerations. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology*, **292**, 18–36.
- Kuo, S.M., MacLean, M.E., McCormick, K. and Wilson, J.X. 2004. Gender and sodium-ascorbate transporter isoforms determine ascorbate concentrations in mice. *Journal of Nutrition*, **134**, 2216–2221.

- Lanfer-Marquez, U.M., Barros, R.M.C. and Sinnecker, P.** 2005. Antioxidant activity of chlorophylls and their derivatives. *Food Research International*, **38**, 885-891.
- Lewis, K.N., Mele, J., Hayes, J.D. and Buffenstein, R.** 2010. Nrf2, a guardian of health span and gatekeeper of species longevity. *Integrative and Comparative Biology*, **50**, 829-843.
- Lescai, F., Marchegiani, F. and Franceschi, C.** 2009. PON1 is a longevity gene: results of a meta-analysis. *Ageing Research Reviews*, **8**, 277-284.
- Liberatore, L., Procida, G., D'Alessandro, N., Cichelli, A.** 2001. Solid-phase extraction and gas chromatographic analysis of phenolic compounds in virgin olive oil. *Food Chemistry*, **73**, 119-124.
- Lin, J.D., Chiou, W.K., Chang, H.Y., Liu, F.H. and Weng, H.F.** 2007. Serum uric acid and leptin levels in metabolic syndrome: a quandary over the role of uric acid. *Metabolism*, **56**, 751-756.
- Liu, Z., Sun, L., Zhu, L., Jia, X., Li, X. and Jia, H.** 2007. Hydroxytyrosol protects retinal pigment epithelial cells from acrolein-induced oxidative stress and mitochondrial dysfunction. *Journal of Neurochemistry*, **103**, 2690-2700.
- Longo, V.D. and Kennedy, B.K.** 2006. Sirtuins in aging and age-related disease. *Cell*, **126**, 257-268.
- Lopez, S., Bermudez, B., Pacheco, Y.M., Villar, J., Abia, R. and Muriana, F.J.** 2008. Distinctive postprandial modulation of beta cell function and insulin sensitivity by dietary fats: monounsaturated compared with saturated fatty acids. *American Journal of Clinical Nutrition*, **88**, 638-644.
- Loru, D., Incani, A., Deiana, M., Corona, G., Atzeri, A., Melis, M.P., Rosa, A. and Dessì, M.A.** 2009. Protective effect of hydroxytyrosol and tyrosol against oxidative stress in kidney cells. *Toxicology and Industrial Health*, **25**, 301-310.
- Lu, S. C.** 1999. Regulation of hepatic glutathione synthesis: current concepts and controversies. *FASEB Journal*, **13**, 1169-1183.
- Lykkesfeldt, J., Hagen, T.M., Vinarsky, V. and Ames, B.N.** 1998. Age-associated decline in ascorbic acid concentration, recycling, and biosynthesis in rat hepatocytes reversal with (R)- α -lipoic acid supplementation. *FASEB Journal*, **12**, 1183-1189.
- Mackness, B., Durrington, P.N. and Mackness, M.I.** 2002. The paraoxonase gene family and coronary heart disease. *Current Opinion in Lipids*, **13**, 357-362.
- Mackness, B., Durrington, P., McElduff, P., Yarnell, J., Azam, N., Watt, M., Mackness, M.** 2003. Low paraoxonase activity predicts coronary events in the Caerphilly Prospective Study. *Circulation*, **107**, 2775-2779.
- Mann, G.E., Rowlands, D.J., Li, F.Y., de W.P. and Siow, R.C.** 2007. Activation of endothelial nitric oxide synthase by dietary isoflavones: role of NO in Nrf2-mediated antioxidant gene expression. *Cardiovascular Research*, **75**, 261-274.
- Mann, G.E., Bonacasa, B., Ishii, T. and Siow, R.C.M.** 2009. Targeting the redox sensitive Nrf2-Keap1 defense pathway in cardiovascular disease:

protection afforded by dietary isoflavones *Current Opinion in Pharmacology*, **9**, 139–145.

- Manna, C., Galletti, P., Cucciolla, V., Montedoro, G. and Zappia, V.** 1999. Olive oil hydroxytyrosol protects human erythrocytes against oxidative damages. *Journal of Nutritional Biochemistry*, **10**, 159-165.
- Manna, C., D'angelo, S., Migliardi, V., Loffredi, E., Mazzoni, O., Morrica, P., Galletti, P. and Zappia, V.** 2002. Protective effect of the phenolic fraction from virgin olive oils against oxidative stress in human cells. *Journal of Agricultural and Food Chemistry*, **50**, 6521-6526.
- Marsilio, V., Campestre, C. and Lanza, B.** 2001. Phenolic compounds change during California-style ripe olive processing. *Food Chemistry*, **74**, 55–60.
- Martín, M.A., Ramos, S., Granado-Serrano, A.B., Rodríguez-Ramiro, I., Trujillo, M., Bravo, L. and Goya, L.** 2010. Hydroxytyrosol induces antioxidant/detoxifiant enzymes and Nrf2 translocation via extracellular regulated kinases and phosphatidylinositol-3-kinase/protein kinase B pathways in HepG2 cells. *Molecular Nutrition and Food Research*, **54**, 956-966.
- Martinez-Gonzalez, J., Rodriguez-Rodriguez, R., Gonzalez- Diez, M., Rodriguez, C., Herrera, M.D. and Ruiz-Gutierrez, V.** 2008. Oleanolic acid induces prostacyclin release in human vascular smooth muscle cells through a cyclooxygenase-2-dependent mechanism. *Journal of Nutrition*, **138**, 443-448.
- Martínez-González, M.A., García-López, M., Bes-Rastrollo, M., Toledo, E., Martínez-Lapiscina, E.H., Delgado-Rodríguez, M., Vazquez, Z., Benito, S. and Beunza, J.J.** 2011. Mediterranean diet and the incidence of cardiovascular disease: A Spanish cohort. *Nutritional Metabolism and Cardiovascular Disease*, **21**, 237-244.
- Mateos, R., Espartero, J.L., Trujillo, M., Rios, J.J., Leon-Camacho, M., Alcudia, F. and Cert, A.** 2001. Determination of phenols, flavones and lignans in virgin olive oils by solid phase extraction and high performance liquid chromatography with diode array ultraviolet detection. *Journal of Agricultural and Food Chemistry*, **49**, 2185-2192.
- Matsugo, S., Kitagawa, T., Minami, S., Esashi, Y., Oomura, Y., Tokumaru, S., Kojo, S., Matsushima, K. and Sasaki, K.** 2000. Age-dependent changes in lipid peroxide levels in peripheral organs, but not in brain, in senescence-accelerated mice. *Neuroscience Letters*, **278**, 105-108.
- May, J.M.** 2000. How does ascorbic acid prevent endothelial dysfunction? *Free Radical Biology and Medicine*, **28**, 1421-1429.
- Meilin, E., Aviram, M. and Hayek, T.** 2010. Paraoxonase 2 (PON2) decreases high glucose-induced macrophage triglycerides (TG) accumulation, via inhibition of NADPH oxidase and DGAT1 activity: studies in PON2-deficient mice. *Atherosclerosis*, **208**, 390-395.
- Mendilaharsu, M., De Stefani, E., Deneo-Pellegrini, H., Carzoglio, J. and Ronco, A.** 1998. Phytosterols and risk of lung cancer: a case-control study in Uruguay. *Lung Cancer*, **21**, 37-45.

- Michels, A.J., Joisher, N. and Hagen, T.M.** 2003. Age-related decline of sodium-dependent ascorbic acid transport in isolated rat hepatocytes. *Archives of Biochemistry and Biophysics*, **410**, 112–120.
- Mills, B.J., Weiss, M.M., Lang, C.A., Liu, M.C. and Ziegler, C.** 2000. Blood glutathione and cysteine changes in cardiovascular disease. *Journal of Laboratory and Clinical Medicine*, **135**, 396-401.
- Montedoro, G., Servili, M., Baldioli, M. and Miniati, E.** 1992. Simple and hydrolyzable phenolic compounds in virgin olive oil. initial characterization of the hydrolyzable fraction. *Journal of Agricultural and Food Chemistry*, **40**, 1577-1580.
- Montedoro, G.F., Servili, M., Baldioli, M., Selvaggini, R., Miniati, E. and Macchioni, A.** 1993. Simple and hydrolyzable compounds in virgin olive oil. Spectroscopic characterizations of the secoiridoid derivatives. *Journal of Agricultural and Food Chemistry*, **41**, 2228-2234.
- Morel, D.W., Hessler, J.R. and Chisolm, G.M.** 1983. Low density lipoprotein cytotoxicity induced by free radical peroxidation of lipid. *Journal of Lipid Research*, **24**, 1070-1076.
- Moreno, J.J.** 2003. Effect of olive oil minor components on oxidative stress and arachidonic acid mobilization and metabolism by macrophages RAW 264.7. *Free Radical Biology and Medicine*, **35**, 1073-1081.
- Morita, T.** 2005. Heme oxygenase and atherosclerosis. *Arteriosclerosis Thrombosis and Vascular Biology*, **25**, 1786-1795.
- Moskovitz, J., Yim, M.B. and Chock, P.B.** 2002. Free radicals and disease. *Archives of Biochemistry and Biophysics*, **397**, 354-359.
- Motilva, M. J., Romero, M. P., Alegre, S. and Girona, J.** 1999. Effect of regulated deficit irrigation in olive oil production and quality. *Acta Horticulturae*, **474**, 377-380.
- Murase, T., Haramizu, S., Ota, N. and Hase, T.** 2009. Suppression of the aging-associated decline in physical performance by a combination of resveratrol intake and habitual exercise in senescence-accelerated mice. *Biogerontology*, **10**, 423-434.
- Nabeshi, H., Oikawa, S., Inoue, S., Nishino, K. and Kawanishi, S.** 2006. Proteomic analysis for protein carbonyl as an indicator of oxidative damage in senescence-accelerated mice. *Free Radical Research*, **40**, 1173-1181.
- Nakanishi, N., Okamoto, M., Yoshida, H., Matsuo, Y., Suzuki, K. and Tatara, K.** 2003. Serum uric acid and risk for development of hypertension and impaired fasting glucose or Type II diabetes in Japanese male office workers. *European Journal of Epidemiology*, **18**, 523-530.
- Napolitano, A., Morales, F., Sacchi, R. and Fogliano, V.** 2008. Relationship between virgin olive oil phenolic compounds and acrylamide formation in fried crisps. *Journal of Agricultural and Food Chemistry*, **56**, 2034–2040.
- Navarro, A., Gomez, C., Sanchez-Pino, M. J. and Gonzalez, H.** 2005. Vitamin E at high doses improves survival, neurological performance, and brain mitochondrial function in aging male mice. *American Journal of Physiology- Regulatory, Integrative and Comprehensive Physiology*, **289**, 1392–1399.

- Newmark, H.L.** 1997. Squalene, olive oil, and cancer risk: a review and hypothesis. *Cancer Biomarkers Epidemiology and Prevention*, **6**, 1101-1103.
- Ng, C.J., Shih, D.M., Hama, S.Y., Villa, N., Navab, M., and Reddy, S.T.** 2005. The paraoxonase gene family and atherosclerosis. *Free Radical Biology and Medicine*, **38**, 153-163,
- Ng, C.J., Bourquard, N., Grijalva, V.** 2006. Paraoxonase-2 deficiency aggravates atherosclerosis in mice despite lower apolipoprotein-B-containing lipoproteins: antiatherogenic role for paraoxonase-2. *Journal of Biological Chemistry*, **281**, 29491-29500.
- Niki E.** 2009. Lipid peroxidation: physiological levels and dual biological effects. *Free Radical Biology and Medicine*, **47**, 469-484.
- Novak, I., Janeiro, P., Seruga, M. and Oliveira-Brett, A.M.** 2008. Ultrasound extracted flavonoids from four varieties of Portuguese red grape skins determined by reverse-phase high-performance liquid chromatography with electrochemical detection. *Analytica Chimica Acta*, **630**, 107-115
- Novak, I., Šeruga, M., Komorsky-Lovric, S.** 2010. Characterisation of catechins in green and black teas using square-wave voltammetry and RP-HPLC-ECD. *Food Chemistry*, **122**, 1283-1289
- Okatani, Y., Wakatsuki, A., Reiter, R.J. and Miyahara, Y.** 2002. Melatonin reduces oxidative damage of neural lipids and proteins in senescence-accelerated mouse. *Neurobiology of Aging*, **23**, 639-644.
- Owen, R.W., Giacosa, A., Hull, W.E., Haubner, R., Würtele, G. and Spiegelhalder, B.** 2000. Olive oil consumption and health: the possible role of antioxidants. *Lancet Oncology*, **1**, 107-112.
- Quiles, J.L., Ramírez-Tortosa, M.C., Gómez, J.A. and José Mataix, J.R.H.** 2002. Role of vitamin E and phenolic compounds in the antioxidant capacity, measured by ESR, of virgin olive, olive and sunflower oils after frying. *Food Chemistry*, **76**, 461-468.
- Quiles, J.L., Farquharson, A.J. and Simpson, D.K.** 2002. Olive oil phenolics: effects on DNA oxidation and redox enzyme mRNA in prostate cells. *British Journal of Nutrition*, **88**, 225-234.
- Packer, L., Weber, S.U. and Rimbach, G.** 2001. Molecular aspects of α -tocotrienol antioxidant action and cell signalling. *Journal of Nutrition*, **131**, 369-373.
- Pallas, M., Verdaguer, E., Tajés, M., Gutierrez-Cuesta, J. and Camin, A.,** 2008. Modulation of sirtuins: new targets for antiageing. *Recent Patents on CNS Drug Discovery*, **3**, 61-69.
- Pallàs, M., Pizarro, J. G., Gutierrez-Cuesta, J., Crespo-Biel, N., Alvira, D., Tajés, M., Yeste-Velasco, M., Folch, J., Canudas, A. M., Sureda, F. X., Ferrer, I. and Camins, A.** 2008. Modulation of Sirt1 expression in different neurodegenerative models and human pathologies *Neuroscience*, **154**, 1388-1397.
- Paniagua, J.A., de la Sacristana, A.G., Sanchez, E., Romero, I., Vidal- Puig, A. and Berral, F.J.** 2007. A MUFA-rich diet improves postprandial glucose, lipid and GLP-1 responses in insulin-resistant subjects. *Journal of American Collaborative Nutrition*, **26**, 434-444.

- Penalvo, J.L. and Nurmi, T.** 2006. Application of coulometric electrode array detection to the analysis of isoflavonoids and lignans. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 1497-1507.
- Peng, Y., Liu, F., Peng, Y. and Ye, J.** 2005. Determination of polyphenols in apple juice and cider by capillary electrophoresis with electrochemical detection. *Food Chemistry*, **92**, 169-175.
- Perona, J.S., Covas, M.I., Fitó, M., Cabello-Moruno, R., Aros, F., Corella, D., Rose, E., Garcia, M., Estruch, R., Martinez-Gonzalez, M.A. and Ruiz-Gutierrez, V.** 2010. Reduction in systemic and VLDL triacylglycerol concentration after a 3-month Mediterranean-style diet in high-cardiovascular-risk subjects. *Journal of Nutritional Biochemistry*, **21**, 892-898.
- Perugini, P., Vettor, M., Rona, C., Troisi, L., Villanova, L., Genta, I., Conti, B. and Pavanetto, F.** 2008. Efficacy of oleuropein against UVB irradiation: preliminary evaluation. *International Journal of Cosmeceutical Science*, **30**, 113-120.
- Petrache, I., Otterbein, L.E., Alam, J., Wiegand, G.W. and Choi, A.M.** 2000. Heme oxygenase-1 inhibits TNF-alpha-induced apoptosis in cultured fibroblasts. *American Journal of Physiology- Lung Cellular and Molecular Physiology*, **278**, 312-319.
- Petursdottir, A.L., Farr, S.A., Morley, J.E., Banks, W.A. and Skuladottir, G.V.** 2008. Effect of Dietary n-3 polyunsaturated fatty acids on brain lipid fatty acid composition, learning ability, and memory of senescence-accelerated mouse. *Journal of Gerontology: Biological Sciences*, **63**, 1153-1160.
- Pinelli, P., Galardi, C., Mulinacci, N., Vinceri, F.F., Cimato, A. and Romani, A.** 2003. Minor polar compound and fatty acid analyses in monocultivar virgin olive oils from Tuscany. *Food Chemistry*, **80**, 331-336.
- Pirisi, F.M., Angioni, A., Cabras, P., Garau, v.L., di Teulada, M.T.S., dos Santos, M.K. and Bandino, G.** 1997. Phenolic compounds in virgin olive oils I. Low-wavelength quantitative determination of complex phenols by high-performance liquid chromatography under isocratic elution. *Journal of Chromatography A*, **768**, 207-213.
- Pitozzi, V., Jacomelli, M., Zaid, M., Luceri, C., Bigagli, E., Lodovici, M., Ghelardini, C., Vivoli, E., Norcini, M., Gianfriddo, M., Esposto, S., Servili, M., Morozzi, G., Baldi, E., Bucherelli, C., Dolara, P. and Giovannelli, L.** 2010. Effects of dietary extra-virgin olive oil on behaviour and brain biochemical parameters in ageing rats. *British Journal of Nutrition*, **103**, 1674-1683.
- Ponnappan, S., Ovaa, H. and Ponnappan, U.** 2007. Lower expression of catalytic and structural subunits of the proteasome contributes to decreased proteolysis in peripheral blood T lymphocytes during aging. *The International Journal of Biochemistry & Cell Biology* **39**, 799-809.
- Prior, R., Wu, X. and Schich, K.** 2005. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, **53**, 4290-4302.
- Quiles, J.L., Ochoa, J.J., Ramirez-Tortosa, C., Battino, M., Huertasa, J.R., Martina, Y. and Mataix, J.** 2004. Dietary fat type (virgin olive vs. sunflower oils) affects age-related changes in DNA double-strand-

- breaks, antioxidant capacity and blood lipids in rats *Experimental Gerontology*, **39**, 1189-1198.
- Ragazzi, E. and Veronese, G.** 1973. Quantitative analysis of phenolic compounds after thin layer chromatographic separation. *Journal of Chromatography*, **77**, 369-375.
- Raitakari, O.T., Salo, P., Gylling, H. and Miettinen, T.A.** 2008. Plant stanol ester consumption and arterial elasticity and endothelial function. *British Journal of Nutrition*, **100**, 603-608.
- Rebrin, I., Zicker, S., Wedekind, K.J., Paetau-Robinson, I., Packer, L. and Sohal, R.S.** 2005. Effect of antioxidant-enriched diets on glutathione redox status in tissue homogenates and mitochondria of the senescence-accelerated mouse. *Free Radical Biology and Medicine*, **39**, 549-557.
- Reddy, S.T., Devarajan, A., Bourquard, N., Shih, D. and Fogelman, A.M.** 2008. Is it just paraoxonase 1 or are other members of the paraoxonase gene family implicated in atherosclerosis? *Current Opinion in Lipidology*, **19**, 405-408.
- Reddy, L.H. and Couvreur, P.** 2009. Squalene: A natural triterpene for use in disease management and therapy. *Advanced Drug Delivery Reviews*, **61**, 1412-1426.
- Rietjens, I.M.C.M., Boersma, M.G., de Haan, L., Spenkelink, B., Awad, H.M., Cnubben, N.H.P., van Zanden, J.J., van der Woude, H., Alink, G.M. and Koeman, J.H.** 2002. The pro-oxidant chemistry of the natural antioxidants vitamin C, vitamin E, carotenoids and flavonoids. *Environmental Toxicology and Pharmacology*, **11**, 321-333.
- Rigacci, S., Guidotti, V., Bucciantina, M., Parri, M., Nediani, C., Cerbai, E., Stefani, M. and Berti, A.** 2010. Oleuropein aglycon prevents cytotoxic amyloid aggregation of human amylin. *Journal of Nutritional Biochemistry*, **21**, 726-735.
- Rimm, E.B., Stampfer, M.J., Ascherio, A., Giovannucchi, E., Colditz, G.A. and Willett, W.C.** 1993. Vitamin E consumption and the risk of coronary heart disease in men. *The New England Journal of Medicine*, **328**, 1450-1456.
- Ristow, M., Zarse, K., Oberbach, A., Klötting, N., Birringer, M., Kiehnopf, M., Stumvoll, M., Kahn, C.R. and Blüher, M.** 2009. Antioxidants prevent health-promoting effects of physical exercise in humans. *Proceedings of the National Academy of Sciences*, **106**, 8665-8670.
- Rodríguez, M.I., Escames, G., López, L.C., López, A., García, J.A., Ortiz, F., Sánchez, V., Romeu, M. and Acuña-Castroviejo, D.** 2008. Improved mitochondrial function and increased lifespan after chronic melatonin treatment in senescent prone mice. *Experimental Gerontology*, **43**, 749-756.
- Rosenblat, M. and Aviram, M.** 2009. Paraoxonases role in the prevention of cardiovascular diseases. *Biofactors*, **35**, 98-104.
- Rosenblat, M., Coleman, R., Reddy, S.T. and Aviram, M.** 2009. Paraoxonase 2 (PON2) attenuates macrophage triglyceride accumulation via inhibition of diacylglycerol acyltransferase 1 (DGAT1), the rate limiting enzyme in triglycerides biosynthesis. *Journal of Lipid Research*, **50**, 870-879.

- Ross, D., Kupa, J.K., Winski, S.L., Beall, H.D., Anwar, A. and Siegel, D. 2000. NAD(P)H:quinone oxidoreductase 1 (NQO1): chemoprotection, bioactivation, gene regulation and genetic polymorphisms. *Chemico-Biological Interactions*, **129**, 77-97.
- Saldeen, T., Li, D. and Mehta, J.L. 1999. Differential effects of α - and γ -tocopherol on low density protein oxidation, superoxide activity, platelet aggregation, and arterial thrombogenesis. *Journal of the American College of Cardiology*, **34**, 1208-1215.
- Santocono, M., Zurria, M., Berrettini, M., Fedeli, D. and Falcioni, G. 2007. Lutein, zeaxanthin and astaxanthin protect against DNA damage in SK-N-SH human neuroblastoma cells induced by reactive nitrogen species. *Journal of Photochemistry and Photobiology B*, **88**, 1-10.
- Sato, E., Oda, N., Ozaki, N., Hashimoto, S., Kurokawa, T. and Ishibashi, S. 1996. Early and transient increase in oxidative stress in the cerebral cortex of senescence-accelerated mouse. *Mechanisms in Ageing and Development*, **86**, 105-114.
- Satoh, A., Yokozawa, T., Cho, E.J., Okamoto, T. and Sei, Y. 2004. Antioxidative effects related to the potential anti-aging properties of the Chinese prescription Kangen-karyu and Carthami Flos in senescence-accelerated mice. *Archives of Gerontology and Geriatrics*, **39**, 69-82.
- Savini, I., Rossi, A., Pierro, C., Avigliano, L. and Catani, M.V. 2008. SVCT1 and SVCT2: key proteins for vitamin C uptake. *Amino Acids*, **34**, 347-355.
- Schaffer, S., Podstawa, M., Visioli, F., Bogani, P., Muller, W.E. and Eckert, G.P. 2007. Hydroxytyrosol-rich olive mill wastewater extract protects brain cells in vitro and ex vivo. *Journal of Agricultural and Food Chemistry*, **55**, 5043-5049.
- Schiborr, C., Eckert, G.P., Weissenberger, J., Müller, W.E., Schwamm, D., Grune, T. and Rimbach, G. 2010. Cardiac oxidative stress and inflammation are similar in SAMP8 and SAMR1 mice and unaltered by curcumin and Ginkgo biloba extract intake. *Current Pharmacology and Biotechnology*, **11**, 861-867.
- Selman, C., McLaren, J.S., Meyer, C., Duncan, J.S., Redman, P., Collins, A.R., Duthie, G.G. and Speakman, J.R. 2006. Life-long vitamin C supplementation in combination with cold exposure does not affect oxidative damage or lifespan in mice, but decreases expression of antioxidant protection genes. *Mechanisms of Ageing and Development*, **127**, 897-904.
- Seppo, L., Jauhiainen, T., Nevala, R., Poussa, T. and Korpela, R. 2007. Plant stanol esters in low-fat milk products lower serum total and LDL cholesterol. *European Journal of Nutrition*, **46**, 111-117.
- Seres, I., Paragh, G., Deschene, E., Fulop, T. and Khalil, A. 2004. Study of factors influencing the decreased HDL associated PON1 activity with aging. *Experimental Gerontology*, **39**, 59-66.
- Serrano, P., Yago, M.A.D., Manas, M., Calpena, R., Mataix, J.A. and Martinez-Victoria, E. 1997. Influence of type of dietary fat (olive and sunflower oil) upon gastric acid secretion and release of gastrin, somatostatin, and peptide YY in Man. *Digestive Diseases and Sciences*, **42**, 626-633.

- Servili, M., Montedoro, G.** 2002. Contribution of phenolic compounds to virgin olive oil quality. *European Journal of Lipid Science and Technology*, **104**, 602–613.
- Servili, M., Selvaggini, R., Esposto, S., Taticchi, A., Montedoro, G. and Morozzi, G.** 2004. Health and sensory properties of virgin olive oil hydrophilic phenols: agronomic and technological aspects of production that affect their occurrence in the oil. *Journal of Chromatography A*, **1054**, 113-127
- Servili, M., Selvaggini, R., Taticchi, A., Esposto, S. and Montedoro, G.** 2003. Volatile compounds and phenolic composition of virgin olive oil: optimization of temperature and time of exposure of olive pastes to air contact during the mechanical extraction process. *Journal of Agricultural and Food Chemistry*, **51**, 7980-7988.
- Shih, P.H., Chan, Y.C., Liao, J.W., Wang, M.F. and Yen, G.C.** 2009. Antioxidant and cognitive promotion effects of anthocyanin-rich mulberry (*Morus atropurpurea* L.) on senescence-accelerated mice and prevention of Alzheimer's disease. *Journal of Nutritional Biochemistry*, **21**, 598-605.
- Simonsen, N.R., Fernandez-Crehuet, N.J. and Martin-Moreno, J.M.** 1998. Tissue stores of individual monounsaturated fatty acids and breast cancer: the EURAMIC study. European Community Multicenter Study on antioxidants, myocardial infarction, and breast cancer. *American Journal of Clinical Nutrition*, **68**,134-141.
- Singleton, V.L., Orthofer, R. and Lamuela-Raventós, R.M.** 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin ciocalteu reagent, *Methods in Enzymology*, **299**, 152-178.
- Škevin, D., Rade, D., Štrucelj, D., Mokrovčak, I., Nederal, S. and Benčić, D.** 2003. The influence of variety and harvest time on the bitterness and phenolic compounds of olive oil. *European Journal of Lipid Science and Technology*, **105**, 536–541.
- Sofi, F., Cesari, F., Abbate, R., Gensini, G.F. and Casini, A.** 2008. Adherence to Mediterranean diet and health status: meta-analysis. *BMJ*. **337**,1344-1350.
- Sohal, R.S.**, 2002. Oxidative stress hypothesis of aging. *Free Radical Biology and Medicine*, **33**, 573-574.
- Solfrizzi, V., Colacicco, A.M., D'Introno, A., Capurso, C., Torres, F. and Rizzo, C.** 2006. Dietary intake of unsaturated fatty acids and agerelated cognitive decline: a 8.5-year follow-up of the Italian longitudinal study on aging. *Neurobiology of Aging*, **27**, 1694-704.
- Solinas, M., Cichelli, A.** 1982. GLC and HPLC evaluation of phenolic compounds in olive oil. *Riv. Soc. Ital. Sci. Aliment.* **11**, 223–230.
- Soriguer, F, Esteva, I, Rojo-Martinez, G, Ruiz de Adana M.S., Dobarganes, M.C. and Garcia-Almeida, J.M.** 2004. Oleic acid from cooking oils is associated with lower insulin resistance in the general population (Pizarra study). *European Journal of Endocrinology*, **150**, 33-39.
- Somova, L.I., Shode, F.O., Rammanan, P. and Nadar, A.** 2003. Antihypertensive, antiatherosclerotic and antioxi- dant activity of triterpenoids isolated from *Olea europaea*, subspecies *africana* leaves. *Journal of Ethnopharmacology*, **84**, 299-305.

- Stadtman, E.R. and Berlett, B.S.** 1997. Reactive oxygen-mediated protein oxidation in aging and disease. *Chemical Research in Toxicology*, **10**, 485-494.
- Stoneham, M., Goldacre, M. and Seagroatt, V.** 2000. Olive oil, diet and colorectal cancer: an ecological study and a hypothesis. *Journal of Epidemiology and Community Health*, **54**, 756-760.
- Strazzullo, P. and Puig, J.G.,** 2007. Uric acid and oxidative stress: relative impact on cardiovascular risk. *Nutrition Metabolism and Cardiovascular Diseases*, **17**, 409-414.
- Svendsen, C. N.** 1993. Multi-electrode array detectors in high-performance liquid chromatography: a new dimension in electrochemical analysis. *Analyst*. **118**, 123-129.
- Syktiotis, G.P., Habeos, I.G., Samuelson, A.W. and Bohmann, D.** 2011. The role of the antioxidant and longevity-promoting Nrf2 pathway in metabolic regulation. *Current Opinion in Clinical Nutrition and Metabolic Care*, **14**, 41-48.
- Tajes, M., Gutierrez-Cuesta, J., Folch, J., Ortuño-Sahagun, D., Verdaguer, E., Jiménez, A., Junyent, F., Lau, A., Camins, A. and Pallàs, M.** 2010. Neuroprotective role of intermittent fasting in senescence-accelerated mice P8 (SAMP8). *Experimental Gerontology*, **45**, 702-710.
- Takeda T., Hosokawa M. and Higuchi K.** 1997. Senescence-accelerated mouse (SAM): a novel murine model of senescence. *Experimental Gerontology*, **32**, 105-109.
- Tafazoli, S., Wright, J.S. and O'Brien, P.J.** 2005. Prooxidant and Antioxidant Activity of Vitamin E Analogues and Troglitazone. *Chemical Research in Toxicology*, **18**, 1567-1574.
- Thomàs-Moyá, E., Gianotti, M., Proenza, A.M., Llado, I.** 2006. The age-related paraoxonase 1 response is altered by long-term caloric restriction in male and female rats. *Journal of Lipid Research*, **47**, 2042-2048.
- Tovar, M.J., Motilva, M.J., Romero, M.P.** 2001. Changes in the phenolic composition of virgin olive oil from young trees (*Olea europaea* L. cv. Arbequina) grown under linear irrigation strategies. *Journal of Agricultural and Food Chemistry*, **49**, 5502-5508.
- Traber, M.G. and Arai, H.** 1999. Molecular mechanisms of vitamin E transport. *Annual Reviews in Nutrition*, **19**, 343-355.
- Trichopoulou, A., Costacou, T., and Bamia, C.** 2003. Adherence to a Mediterranean diet and survival in a Greek population. *New England Journal of Medicine*, **348**, 2599-2608.
- Tripoli, E., Giammanco, M., Tabacchi, G., Di Majo, D., Giammanco, S. and La Guardia, M.** 2005. The phenolic compounds of olive oil: structure, biological activity and beneficial effects on human health. *Nutrition Research Reviews*, **18**, 98-112.
- Trumbo, P.R. and Ellwood, K.C.** 2006. Lutein and zeaxanthin intakes and risk of age-related macular degeneration and cataracts: an evaluation using the Food and Drug Administration's evidence-based review system for health claims. *American Journal of Clinical Nutrition*, **84**, 971-974.

- Tsimidou, M., Papadopoulos, G., and Boskou, D.** 1992. Determination of phenolic compounds in virgin olive oil by reversed-phase HPLC with emphasis on UV detection. *Food Chemistry*, **44**, 53-60.
- Tzonou, A., Hsieh, C. and Polychronopoulou, A.** 1993. Diet and ovarian cancer: a case-control study in Greece. *International Journal of Cancer*, **55**, 411-414.
- Unno, K., Takabayashi, F., Kishido, T. and Oku, N.** 2004. Suppressive effect of green tea catechins on morphologic and functional regression of the brain in aged mice with accelerated senescence (SAMP10). *Experimental Gerontology*, **39**, 1027-1034.
- Valko, M., Leibfritz D., Moncol J., Cronin M.T., Mazur M. and Telser, J.** 2007. Free radicals and antioxidants in normal physiological functions and human disease. *International Journal of Biochemical Cell Biology*, **39**, 44-84.
- Vauzour, D., Vafeiadou, K., Corona, G., Pollard, S.E., Tzounis, X. and Spencer, J.P.** 2007. Champagne wine polyphenols protect primary cortical neurons against peroxynitrite-induced injury. *Journal of Agricultural and Food Chemistry*, **55**, 2854-2860.
- Vanbeneden, N., Delvaux, F. and Delvaux, F.R.** 2006. Determination of hydroxycinnamic acids and volatile phenols in wort and beer by isocratic high-performance liquid chromatography using electrochemical detection. *Journal of Chromatography A*, **1136**, 237-242.
- van der Loo, B., Labugger, R., Aebischer, C.P., Skepper, J.N., Bachschmid, M., Spitzer, V., Kilo, J., Altwegg, L., Ullrich, V. and Lüscher, T.F.** 2002. Cardiovascular Aging Is Associated With Vitamin E Increase. *Circulation*, **105**, 1635-1638.
- van Rensburg, S.J., Daniels, W.M., van Zyl, J.M. and Taljaard, J.J.** 2000. A comparative study of the effects of cholesterol, beta-sitosterol, betasitosterol glucoside, dehydroepiandrosterone sulphate and melatonin on in vitro lipid peroxidation. *Metabolic Brain Disease*, **15**, 257-265.
- Verhoef, P., Stampfer, M.J., Buring, J.E., Gaziano, J.M., Allen, R.H. and Stabler, S.P.** 1996. Homocysteine metabolism and risk of myocardial infarction: relation with vitamins B6, B12, and folate. *American Journal of Epidemiology*, **143**, 845-859.
- Visioli, F., Bellosta, S. and Galli, C.** 1998a. Oleuropein, the bitter principle of olives, enhances nitric oxide production by mouse macrophages. *Life Sciences*, **62**, 541-546.
- Visioli, F., Bellomo, G. and Galli, C.** 1998b. Free radical-scavenging properties of olive oil polyphenols. *Biochemistry Biophysics Research Communications*, **247**, 60-64.
- Visioli, F. and Galli, C.** 2000. Olive oil: more than just oleic acid. *American Journal of Clinical Nutrition*, **72**, 853-856.
- Visioli, F., Poli, A. and Galli, C.** 2002. Antioxidant and other biological activities of phenols from olives and olive oil. *Medicinal Research Reviews*, **22**, 65-75.

- Visioli, F., Bellomo, G., Galli, C. and Caruso, D. 2002. Biological activities and metabolic fate of olive oil phenols. *European Journal of Lipid Science and Technology*, **104**, 677–684.
- Viteri, G., Carrard, G., Birlouez-Aragon, I., Silva, E. and Friguet, B. 2004. Age-dependent protein modifications and declining proteasome activity in the human lens. *Archives of Biochemistry and Biophysics*, **427**, 197–203.
- Wilkinson, J. and Clapper, M.L. 1997. Detoxification enzymes and chemoprevention. *Proceedings of the Society for Experimental Biology and Medicine*, **216**, 192–200.
- Wu, T., Guan, Y. and Ye, J. 2007. Determination of flavonoids and ascorbic acid in grapefruit peel and juice by capillary electrophoresis with electrochemical detection. *Food Chemistry*, **100**, 1573–1579.
- Xu, C., Yong-Tao Li, C. and Tony Kong, A.N. 2005. Induction of Phase I, II and III Drug Metabolism/Transport by Xenobiotics. *Archives of Pharmacological Research*, **28**, 249–268.
- Xu, X., Yu, L. and Chen, G. 2006. Determination of flavonoids in *Portulaca oleracea* L. by capillary electrophoresis with electrochemical detection. *Journal of Pharmacology and Biomedical Analysis*, **41**, 493–499.
- Yamaki, T., Yano, T., Satoh, H. and Endo, T. 2002. High oleic acid oil suppresses lung tumorigenesis in mice through the modulation of extracellular signal-regulated kinase cascade. *Lipids*, **37**, 783–788.
- Yasui, F., Ishibashi, M., Matsugo, S., Kojo, S., Oomura, Y. and Sasaki, K. 2003. Brain lipid hydroperoxide level increases in senescence-accelerated mice at an early age. *Neuroscience Letters*, **350**, 66–68.
- Ye, X., Meeker, H.C., Kozlowski, P.B., Wegiel, J., Wang, K.C., Imaki, H. and Carp, R.I. 2004. Pathological changes in the liver of a senescence accelerated mouse strain (SAMP8): A mouse model for the study of liver diseases. *Histology and Histopathology*, **19**, 1141–1151.
- Young, J., Wahle, K.W.J. and Boyle, S.P. 2008. Cytoprotective effects of phenolic antioxidants and essential fatty acids in human blood monocyte and neuroblastoma cell lines: Surrogates for neurological damage in vivo. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, **78**, 45–59.
- Yu, B.P. and Chung, H.Y. 2006. The inflammatory process in aging. *Reviews in Clinical Gerontology*, **16**, 179–187.
- Zeng, B.Y., Medhurst, A.D., Jackson, M., Rose, S. and Jenner, P. 2005. Proteasomal activity in brain differs between species and brain regions and changes with age. *Mechanisms of Ageing and Development*, **126**, 760–766.
- Zhang, X., Cao, J. and Zhong, L. 2009. Hydroxytyrosol inhibits pro-inflammatory cytokines, iNOS, and COX-2 expression in human monocytic cells. *Naunyn-Schmied Archives of Pharmacology*, **379**, 581–586.
- Zhu, H., Itoh, K., Yamamoto, M., Zweier, J.L., and Li, Y. 2005. Role of Nrf2 signaling in regulation of antioxidants and phase 2 enzymes in cardiac fibroblasts: Protection against reactive oxygen and nitrogen species-induced cell injury. *FEBS Letters*, **579**, 3029–3036.
- Zhu, B.H., Ueno, M. and Matsushita, T. 2001. Effects of aging and blood pressure on the structure of the thoracic aorta in SAM mice: a model of age-

associated degenerative vascular changes. *Experimental Gerontology*, **36**, 111-124.

Zrelli, H, Matsuoka, M., Kitazaki, S., Araki, M., Kusunoki, M., Zarrouk, M. and Miyazaki, H. 2011. Hydroxytyrosol induces proliferation and cytoprotection against oxidative injury in vascular endothelial cells: role of Nrf2 activation and HO-1 induction. *Journal of Agricultural and Food Chemistry*, **59**, 4473-4482.

APPENDICES

APPENDIX A: Chromatograms

APPENDIX B: Calibration curves

APPENDIX C: List of olive oils that are analysed in the study

APPENDIX D: Statistical values

APPENDIX A: Chromatograms




Figure A.1: Representative chromatogram of the standards 1) hydroxytyrosol 2) tyrosol 3) caffeic acid 4) vanillic acid 5) ferulic acid 6) p-coumaric acid 7)oleuropein 8) pinoresinol IS: internal standard.




Figure A.2: Representative chromatogram of the olive oil extract1) hydroxytyrosol 2) tyrosol 3) caffeic acid 4) vanillic acid 5) ferulic acid 6) p-coumaric acid 7) oleuropein 8) pinoresinol IS: internal standard.




Figure A.3: Representative chromatogram of the blank olive oil extract.




Figure A.4: Representative chromatogram of α - and γ -tocopherol.

APPENDIX B: Calibration Curves




Figure B.1: Calibration curve for FRAP assay.




Figure B.2: Calibration curve for TEAC assay.




Figure B.3: Calibration curve for the determination of total phenolics.




Figure B.4: Calibration curve for tyrosol.




Figure B.5: Calibration curve for hydroxytyrosol.




Figure B.6: Calibration curve for vanillic acid.




Figure B.7: Calibration curve for caffeic acid.




Figure B.8: Calibration curve for p-coumaric acid.




Figure B.9: Calibration curve for ferulic acid.




Figure B.10: Calibration curve for oleuropein.




Figure B.11: Calibration curve for pinoresinol.

APPENDIX C:

Table C.1: The list of olive oil samples that are analyzed in the study.

Sample	Country	Region	Olive variety	Harvest year	Extraction	Type
1	Spain	n.a.	Lechin	2007	Cold pressed	Extra virgin
2	Spain	n.a.	Picual	2007	Cold pressed	Extra virgin
3	Spain	n.a.	Hojiblanca	2007	Cold pressed	Extra virgin
4	Spain	n.a.	Arbequina	2007	Cold pressed	Extra virgin
5	Spain	Andalusia	Picual	2008	Cold pressed	Extra virgin
6	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
7	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
8	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
9	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
10	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
11	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
12	Spain	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
13	Turkey	Ezine	n.a.	2008	Cold pressed	Extra virgin
14	Turkey	Zeytindağ	n.a.	2008	Cold pressed	Extra virgin
15	Turkey	Aydin	Memecik	2007	Cold pressed	Extra virgin
16	Turkey	Hatay	Saurani	n.a.	Cold pressed	Extra virgin
17	Turkey	Hatay	Hasebi	n.a.	Cold pressed	Extra virgin
18	Turkey	Hatay	Halhali	n.a.	Cold pressed	Extra virgin
19	Turkey	Hatay	Multivariety	n.a.	Cold pressed	Extra virgin
20	Turkey	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
21	Turkey	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
22	Turkey	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
23	Turkey	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
24	Turkey	Ayvalik	Ayvalik	2007	Cold pressed	Extra virgin
25	Italy	Sicily	Tonda iblea	2008	Cold pressed	Extra virgin
26	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
27	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
28	Italy	n.a.	n.a.	2007	Cold pressed	Extra virgin
29	Italy	Sicily	Cerasuola,Nocellara,Biancolilla	2007	Cold pressed	Extra virgin

Table C.1 (continued): The list of olive oil samples that are analyzed in the study.

Sample	Country	Region	Olive variety	Harvest year	Extraction	Type
30	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
31	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
32	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
33	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
34	Italy	n.a.	n.a.	n.a.	Cold pressed	Extra virgin
35	Greece	Lakonia	Koroneiki,Athinoleia	2007	Cold pressed	Extra virgin
36	Greece	Kalamata	Koroneiki	2007	Cold pressed	Extra virgin
37	Greece	Crete	Koroneiki	2007	Cold pressed	Extra virgin
38	Greece	Sitia, Crete	Koroneiki	2007	Cold pressed	Extra virgin
39	Greece	Crete	Koroneiki	2007	Cold pressed	Extra virgin
40	Greece	Crete	Koroneiki	2007	Cold pressed	Extra virgin
41	Greece	Mani	Koroneiki	2007	Cold pressed	Extra virgin
42	Greece	Kolymvari, Crete	Koroneiki	n.a.	Cold pressed	Extra virgin
43	France	Ollioules	Brun	2008	Cold pressed	Extra virgin
44	France	Ollioules	Cayon	2008	Cold pressed	Extra virgin
45	France	Ollioules	Multivariety	2008	Cold pressed	Extra virgin
46	France	n.a.	n.a.	2008	Cold pressed	Extra virgin
47	Portugal	Alentejo	80% Arbequina, 20%Cobrançosa	2007	Cold pressed	Extra virgin
48	Portugal	Alentejo	80% Arbequina, 20%Cobrançosa	2007	Cold pressed	Extra virgin
49	Portugal	n.a.	n.a.	2007	Cold pressed	Extra virgin
50	Portugal	n.a.	n.a.	2007	Cold pressed	Extra virgin
51	Australia	Kiewa River	Lechino	2008	Cold pressed	Extra virgin
52	Australia	Frankland River	Lechino	2007	Cold pressed	Extra virgin
53	Australia	Frankland River	Frantoio	2007	Cold pressed	Extra virgin
54	USA	California	Manzanillo, Mission	2007	Cold pressed	Extra virgin
55	USA	California	Lechino	2007	Cold pressed	Extra virgin
56	USA	California	Frantoio	2007	Cold pressed	Extra virgin
57	South Africa	Morgenster	Frantoio, leccino, coratina	2007	Cold pressed	Extra virgin

Abbreviations: n.a; not available

APPENDIX D: Statistical values

Table D.1: p values of t-test for the analysed parameters in serum of mice

ANALYSED PARAMETER	SAMR1 LP vs SAMP8 LP	SAMP8 LP vs SAMP8 HP
Triglyceride	0.29	0.07
Total cholesterol	0.002	0.23
HDL	0.13	0.84
PON1 activity	0.0001	0.024

Table D.2: p values of t-test for the analysed parameters in heart of mice

ANALYSED PARAMETER	SAMR1 LP vs SAMP8 LP	SAMP8 LP vs SAMP8 HP
TBARS	0.013	0.0001
Protein carbonyl	0.031	0.004
Proteasomal activity		
β -1 subunit	0.173	0.15
β -2 subunit	0.06	0.06
β -5 subunit	0.357	0.35
GSH	0.13	0.088
Cysteine	0.03	0.03
Methionine	0.17	0.10
Uric acid	0.021	0.027
Ascorbic acid	0.12	0.10
Nrf-2	0.76	0.036
GST	0.005	0.10
γ -GCS	0.66	0.06
NQO-1	0.001	0.028
HO-1	0.003	0.86
PON-2	1.00	0.004
SIRT-1	0.029	0.006

Table D.3: p values of t-test for the analysed parameters in liver of mice

ANALYSED PARAMETER	SAMR1 LP vs SAMP8 LP	SAMP8 LP vs SAMP8 HP
TBARS	0.002	0.66
Protein carbonyl	0.016	0.008
Proteasomal activity		
β-1 subunit	0.63	0.98
β-2 subunit	0.87	0.26
β-5 subunit	0.02	0.16
GSH	0.63	0.33
GSSG	0.10	0.56
Cysteine	0.005	0.46
Methionine	0.86	0.51
Uric acid	0.20	0.16
Ascorbic acid	0.003	0.76
SVCT1	0.029	0.71
SVCT2	0.38	0.29
GLUOX	0.77	0.0001
Vitamin E	0.0001	0.0001
CYP3A3	0.28	0.81
TTP	0.32	0.69
LPL	0.08	0.53
LRP1	0.77	0.92
SRB1	0.76	0.09
MDR-1	0.0001	0.11
LDLR	0.36	0.72
ABCA1	0.22	0.17
Nrf-2	0.09	0.71
GST	0.16	0.12
GST (protein level)	0.84	0.15
γ-GCS	0.22	0.33
γ-GCS (protein level)	0.49	0.40
NQO-1	0.069	0.53
HO-1	0.24	0.65
HO-1 (protein level)	0.01	0.87
PON-1	0.47	1.00
SIRT-1	0.35	0.33
SAP	0.016	0.57
CRP	0.004	0.30

CURRICULUM VITAE

Candidate's full name: Banu BAYRAM

Place and date of birth: Istanbul 14.06.1980

Universities and Colleges attended:

Private Cultur Science High School 1995-1998

Istanbul University 1998- 2002

Istanbul Technical University 2002- 2005

Istanbul Technical University 2005- 2011

Publications:

Papers

Bayram, B., Ozcelik, B., Grimm, S., Roeder, T., Schrader, C., Ernst, I.M.A., Wagner, A.E., Grune, T., Frank, J., Rimbach, G. 2011. A diet rich in olive oil phenolics reduces oxidative stress in the heart of SAMP8 mice by induction of Nrf2-dependent gene expression. *Rejuvenation Research* (Accepted).

Bayram, B., Ozcelik, B. 2008. Trans Yağ Asitleri ve Analiz Yöntemleri, *Dünya Gıda*, July, p 82-87

Bayram, B., Ozcelik, B. 2008. Flavanoidler ve Flavonoidlerce Zengin Besinlerin Kalp Damar Hastalıkları Üzerine Etkileri: *In vivo* ve *In vitro* Çalışmalar, *Hasad Gıda*, July-August, p 38-43

Book Chapters

Özçelik, B., Uygun U., **Bayram, B.** 2010. Seafoods and Environmental Contaminants. In "Handbook of Seafood Quality, Safety and Health Applications" (Edts:C. Alasalvar, Fereidoon Shahidi, Kazuo Miyashita and Udaya Wanasundara). Wiley-Blackwell Publishing, Chapter 25, p 303-317, UK. Print ISBN: 9781405180702, Online ISBN:9781444325546

International Conference Presentations

Bayram, B., Frank, J., Özçelik, B., Grimm, S., Grune, T., Rimbach, G. 2011. Effect of a Diet Rich in Olive Oil Phenolics on Age-Related Changes in Heart Oxidant/Antioxidant Status and Nrf2 Dependent Gene Expression. 5th International Congress on Polyphenols and Health, 17-21 October, Barcelona, Spain (poster presentation)

Bayram, B., Frank, J., Özçelik, B., Rimbach, G. 2011. Increased Expression of Transcription Factor Nrf2 and its Target Genes with a Diet Rich in Olive Oil Phenolics in the Heart Tissue of Ageing Model Senescence Accelerated Mouse

Prone 8 (SAMP8) Mice, Eurofed Lipid Congress, 18-21 September, Rotterdam, The Netherlands, p146 (poster presentation).

Bayram, B., Özçelik, B. 2011. DNA Based Methods as Powerful Techniques for the Authentication of Olive Oil. Eurofed Lipid Congress, 18-21 September, Rotterdam, The Netherlands, p149 (poster presentation).

Bayram, B., Frank, J., Özçelik, B., Grimm, S., Grune, T., Rimbach, G. 2011. A Diet Rich in Olive Oil Phenolics Affects Age-Related Changes in Heart Oxidant/Antioxidant Status, Nrf2 Dependent Gene Expression and SIRT1 Status, SFRR Meeting 2011, September 7-10, Istanbul, Turkey, p83 (poster presentation)

Bayram, B., Frank, J., Esatbeyoglu, T., Özçelik, B., Rimbach, G. 2011. A Validated Liquid Chromatography Method with a Sensitive Electrochemical Detector for the Analysis of Selected Phenolic Compounds in Olive Oil, AOCS World Conference on Oilseed Processing, Fats and Oil Processing, Biofuels and Applications, 21-23 June, Izmir, Turkey (poster presentation)

Bayram, B., Özçelik, B., Tatlı, A. 2010. Polyphenolic Olive Mill Water Extract: A High Added Value By-Product of Olive Oil Industry, 9th EuroFed Lipid Congress, 21-24 November, München, Germany (poster presentation)

Bayram, B., Özçelik, B. 2008. Nanotoxicity and Related Regulations in Food Industry, Nanoscience and Nanotechnology Conference, 9-13 June, Istanbul, Turkey (poster presentation)

Bayram, B., Özçelik, B. 2008. Production of Food Grade Carotenoids From Microorganisms, 2nd COST 928 Annual Meeting, 15-17 October, Istanbul, Turkey (poster presentation)

Bayram, B., Heperkan, D. 2006. Exopolysaccharide Production of Lactic Acid Bacteria Isolated From Boza. FoodMicro, The 20th International ICFMH Symposium, August 29th –September 2nd, Bologna, Italy, p349 (poster presentation)

Bayram, B., Heperkan, D. 2006. Enzyme Profile of Lactic Acid Bacteria Isolated From Boza. IFT annual Meeting and Food Expo, 24-28 June, Orlando, Florida, USA (poster presentation)

Bayram, B., Satır Toklu, G., Ballıkaya, S. 2004. Application of Genetic Techniques in Dairy Industry, International Dairy Symposium, 24-28 May, Isparta, Turkey, p334 (1st best poster award, poster presentation)

National Conference Presentations

Bayram, B., Ozcelik, B. 2010. Su Ürünleri Tüketimindeki Tehlike: Çevresel Kontaminantlar. 2nd National Food Safty Congress, 9-10 December, Istanbul, Turkey (poster presentation)

Ozcelik, B., Aydın, N.N., **Bayram, B.** 2010. Türkiye’de Tüketicilerin Gıda Katkı Maddelerine Karşı Davranışları. 2nd National Food Safty Congress, 9-10 December, Istanbul, Turkey (poster presentation)

Ozcelik, B., **Bayram, B.** 2008. Sarapların Otantifikasyonunda Kullanılan Yöntemler, National Wine and Viniculture Symposium, 6-8 November, Denizli, Turkey (poster presentation)

Bayram, B., Heperkan, D. 2006. Bozadan İzole Edilen Laktik Asit Bakterilerinin Ekzopolisakkarit Oluşturma Kapasitelerinin Belirlenmesi, 9th Food Congress, 24-26 May, Bolu, Turkey, p591 (poster presentation)

Bayram, B., Ozcelik, B. 2003. Antioksidanların Mikotoksinler Üzerine Etkileri, 1st National Mycotoxin Symposium, 18-19 September, Istanbul, Turkey, p198 (poster presentation)